

STAT 535 – Homework 2 Solutions

1. The Beta(1, 2) prior implies Kimya believes it's somewhat unlikely the store is open. The Beta(0.5, 1) prior implies Fernando believes STRONGLY that the store is NOT open. The Beta(3, 10) prior implies Ciara believes that the store is not open, but is not as certain as Fernando. The Beta(2, 0.1) prior implies Taylor believes STRONGLY that the store IS open.




Figure 1: Problem 4.4: Beta priors.

2. (a) $y = 8, n = 10$ (b) $y = 3, n = 13$ (c) $y = 2, n = 16$ (d) $y = 7, n = 10$ (e) $y = 3, n = 6$ (f) $y = 29, n = 31$




Figure 2: Problem 4.10: Beta priors, scaled likelihoods, posteriors.

3. (a) Beta(2, 3) prior with $y = 3, n = 5 \Rightarrow \text{Beta}(2+3, 3-5+3) \Rightarrow \text{Beta}(5, 5)$ posterior.

(b) Beta(5,5) prior with $y = 1, n = 5 \Rightarrow$ Beta(6,9) posterior.
(c) Beta(6,9) prior with $y = 1, n = 5 \Rightarrow$ Beta(7,13) posterior.
(d) Beta(7,13) prior with $y = 2, n = 5 \Rightarrow$ Beta(9,16) posterior.

4. > bechdel %>% filter(year==1980) %>% tabyl(binary)
binary n percent
FAIL 10 0.7142857
PASS 4 0.2857143
> summarize_beta_binomial(alpha = 1, beta = 1, y = 4, n = 14)
model alpha beta mean mode var sd
1 prior 1 0.5000 NaN 0.08333333 0.2886751
2 posterior 5 11 0.3125 0.2857143 0.01263787 0.1124183
> bechdel %>% filter(year==1990) %>% tabyl(binary)
binary n percent
FAIL 9 0.6
PASS 6 0.4
> summarize_beta_binomial(alpha = 5, beta = 11, y = 6, n = 15)
model alpha beta mean mode var sd
1 prior 5 11 0.3125000 0.2857143 0.012637868 0.11241827
2 posterior 11 20 0.3548387 0.3448276 0.007154006 0.08458136
>
> bechdel %>% filter(year==2000) %>% tabyl(binary)
binary n percent
FAIL 34 0.5396825
PASS 29 0.4603175
> # Next, 2000 data:
> summarize_beta_binomial(alpha = 11, beta = 20, y = 29, n = 63)
model alpha beta mean mode var sd
1 prior 11 20 0.3548387 0.3448276 0.007154006 0.08458136
2 posterior 40 54 0.4255319 0.4239130 0.002573205 0.05072677

> # All data at once:
> bechdel %>% filter(year==1980|year==1990|year==2000) %>% tabyl(binary)
binary n percent
FAIL 53 0.576087
PASS 39 0.423913
>
> summarize_beta_binomial(alpha = 1, beta = 1, y = 39, n = 92)
model alpha beta mean mode var sd
1 prior 1 1 0.5000000 NaN 0.083333333 0.28867513
2 posterior 40 54 0.4255319 0.423913 0.002573205 0.05072677

5. (a) The R code is given on the course web page. The posterior mean and median for σ^2 is around 228.2. The posterior mean and median for μ is around 225.2. A 95% HPD credible interval for σ^2 is around (214.95, 241.9). A 95% HPD credible interval for μ is around (217.1, 233.51).

- (b) The R code is given on the course web page. The posterior mean and median for μ is around 223.2. A 95% HPD credible interval for μ is around (216.8, 229.7).
- (c) The point estimate for μ is slightly different in the two cases. But the most notable difference is that the credible interval for μ is narrower when σ^2 is assumed known, which makes sense, since there is less randomness in the model.
6. (a) We need $s/r = 5$ and $\sqrt{\frac{s}{r^2}} = 0.25$. Set $s = 5r$ and substitute into the second equation, and solving these equations yields $s = 400, r = 80$.




Figure 3: Problem 5.5: prior.

- (b) The R code `1-pgamma(10, shape=400, rate=80)` shows that this probability is approximately 0.
7. (a), (b)




Figure 4: Problem 5.6: prior, likelihood, posterior.

(c)

```
> summarize_gamma_poisson(shape = 400, rate = 80, sum_y = 49, n = 6)
      model shape rate    mean     mode     var     sd
1   prior    400    80 5.000000 4.987500 0.06250000 0.2500000
2 posterior   449    86 5.22093 5.209302 0.06070849 0.2463909
```

(d) Seeing the data has slightly increased my estimate of the Poisson mean λ , from a prior estimate of 5 to a posterior estimate of around 5.2.

8. (a)

```
> control_subjects <- football %>%
+   filter(group == "control")
> control_subjects %>%
+   summarize(mean(volume))
  mean(volume)
1           7.6026
```

(b)

```
> summarize_normal_normal(mean = 6.5, sd = 0.4, sigma = 0.5,
+                           y_bar = 7.6026, n = 25)
      model     mean     mode     var     sd
1   prior 6.500000 6.500000 0.16000000 0.40000000
2 posterior 7.537741 7.537741 0.009411765 0.09701425
```

(c)




Figure 5: Problem 5.12: prior, likelihood, posterior.

Seeing the data has substantially increased my estimate of the normal mean μ , from a prior estimate of 6.5 to a posterior estimate of around 7.5.

9. (a) The posterior is proportional to

$$\begin{aligned} L(\theta|y)p(\theta) &\propto \theta(1-\theta)^{y-1}\theta^{\alpha-1}(1-\theta)^{\beta-1} \\ &= \theta^{(\alpha+1)-1}(1-\theta)^{(y+\beta-1)-1} \end{aligned}$$

This posterior is a Beta($\alpha + 1, y + \beta - 1$) distribution.

(b) Yes, since the prior and posterior are both Beta distributions, the Beta is a conjugate prior.

10. (a) The posterior is proportional to

$$\begin{aligned} L(\lambda|y_1, \dots, y_n)p(\lambda) &\propto [\lambda^{y_1} e^{-\lambda}] \cdots [\lambda^{y_n} e^{-\lambda}] e^{-\lambda} \\ &= [\lambda^{\sum_{i=1}^n y_i}] e^{-(n+1)\lambda} \end{aligned}$$

This posterior is a gamma($\sum_{i=1}^n y_i + 1, n + 1$) distribution.

(b) For this data set, $\sum_{i=1}^n y_i = 33$ and $n = 15$, so the posterior is gamma(34, 16). R code:

```
# sum of Y's = 33, n = 15
# Posterior median:
qgamma(.5, shape=33+1, rate=15+1)
# Posterior mean:
34/16
# quantile-based interval:
qgamma(c(.025, .975), shape=33+1, rate=15+1)
# HPD interval:
hpd(qgamma, shape=33+1, rate=15+1)
```

The posterior mean is $34/16 = 2.125$. The 95% quantile-based credible interval is (1.47, 2.90). The 95% HPD credible interval is (1.44, 2.85).