

STAT 515 – Sections 5.6 Supplement

Brian Habing – University of South Carolina
Last Updated: February 2, 2017

S5.6 – Relating the Poisson and Exponential

Section 4.5 introduces the Poisson random variable with 4 basic characteristics. We say that something that follows these rules follows a Poisson process.

- 1) The experiment consists of counting the number of times (let's use Y this time) that a certain event occurs during some given amount of time (or area, volume, weight, distance, or any other unit of measurement).
- 2) The probability that an event occurs in any unit of measurement is the same for all units of that size.
- 3) The number of events that occur in one unit of measurement is independent of the number that occur in other units.
- 4) The mean number of events that occur in one unit of measurement is denoted by the Greek letter lambda (λ).

As given on page 215 the probability mass function for a Poisson random variable is $P[Y = y] = \frac{\lambda^y e^{-\lambda}}{y!}$ for $x=0, 1, 2, \dots$.

Imagine that we have something that follows a Poisson process with parameter λ /unit, but we want to know about the random variable X =amount of time until the first event. One way to approach this is to try and figure out the cumulative distribution function for X .

$$\begin{aligned}P[X \leq x] &= P[1^{\text{st}} \text{ event happens before the } x^{\text{th}} \text{ unit of time}] \\&= 1 - P[1^{\text{st}} \text{ event happens after the } x^{\text{th}} \text{ unit of time}] \\&= 1 - P[\text{there are 0 events before time } x]\end{aligned}$$

The probability of having no failures before a given time is a Poisson random variable! The only complication is adjusting the rate to be over x units of time instead of 1 unit.

$$\frac{\lambda}{(1 \text{ unit of time})} = \frac{x}{x} \cdot \frac{\lambda}{(1 \text{ unit of time})} = \frac{x\lambda}{(x \text{ units of time})}$$

So, plugging in the probability of seeing 0 failures for a Poisson distribution with parameter $x\lambda$ gives:

$$\begin{aligned}P[X \leq x] &= 1 - P[\text{there are 0 failures before time } x] \\&= 1 - \frac{(x\lambda)^0 e^{-x\lambda}}{0!} \\&= 1 - e^{-x\lambda}\end{aligned}$$

This also gives

$$P[X \geq x] = P[X > x] = 1 - P[X \leq x] = e^{-x\lambda}$$

Compare this to the formula for Figure 5.28 in the middle of page 268. If we set $\lambda=1/\theta$ this is exactly the same as an exponential distribution!

So there are two ways to write out (or “parameterize”) an exponential distribution:

Parameterization 1: If λ is the parameter for a Poisson process (the number of events expected in each unit of time), then:

$$f(y) = \lambda e^{-x\lambda} \quad P[X \leq x] = 1 - e^{-x\lambda} \quad \mu = \frac{1}{\lambda} \quad \sigma = 1/\lambda$$

This parameterization with lambda is the one that R uses (lambda is the “rate parameter”).

Parameterization 2: If θ is the expected amount of time until the first occurrence, then:

$$f(y) = \frac{1}{\theta} e^{-x/\theta} \quad P[X \leq x] = 1 - e^{-x/\theta} \quad \mu = \theta \quad \sigma = \theta$$

In R, you need to use $1/\theta$ as the rate parameter. To answer example 5.14 in R we would use $1 - P[X \leq 5] = 1 - \text{pexp}(5, 1/2)$ where the $1/2$ is $1/\theta$.

The particular parameterization you should use depends on what information you were given. Were you told how many events were expected to occur in a unit of time (λ) or how long you expected to wait until the first event (θ)?

You can also use the relationship $\lambda=1/\theta$ to switch between the two ways of looking at it.

Example 5.15 (page 269) continued

- e) What is the λ parameter of the Poisson process that the magnetron tubes life-spans follow?
- f) What is the expected number of tube failures in one year?
- g) Using R, what is the probability of observing two or fewer failures in one year?

Solution: $\theta = 6.25$ years is the average lifespan, so the parameter $\lambda = 1/\theta = 1/6.25 \text{ years} = 0.16/\text{year}$ since λ is the mean of the Poisson distribution. The expected number of failures in one year is 0.16. Using R to find $P[Y \leq 2]$ for a Poisson 0.16, would be `ppois(2, .16)` which is approximately 99.94%

Example 4.14 (page 215) continued:

- e) What is the expected amount of time until the first whale is observed?
- f) What is the probability that the first whale will be observed by half-way through the week?

Solution: This is a Poisson process with $\lambda = 2.6/\text{week}$. The expected time until the first whale is observed is the expected value of the exponential distribution with this parameter, so $1/\lambda = 1/(2.6/\text{week}) = \text{week}/0.385 \approx 0.385$ weeks which is around 2.7 days. The probability that the first whale will be observed by half-way through the week is $P[X \leq 0.5]$ for the exponential distribution $= 1 - e^{-\lambda x} = 1 - e^{-2.6(0.5)} = 1 - e^{-1.3} \approx 0.727$. This could also be found using `pexp(0.5, 2.6)` in R.