

Hypothesis testing

Dr. Yen-Yi Ho

Department of Statistics, University of South Carolina

Hypothesis testing

Recall the one-sample normal model

$$Y_1, \dots, Y_n \stackrel{iid}{\sim} N(\mu, \sigma^2).$$

We may perform a t-test to determine whether μ is equal to some specified value μ_0 .

The **test statistic** gives information about whether $\mu = \mu_0$ is plausible:

$$t^* = \frac{\bar{Y} - \mu_0}{s/\sqrt{n}}.$$

If $\mu = \mu_0$ is true, then $t^* \sim t_{n-1}$.

Hypothesis testing

Test of hypothesis: answer a **yes**, or **no** question regarding a population parameter.

Rationale: Since \bar{y} is our best estimate of the unknown μ , $\bar{y} - \mu_0$ will be small if $\mu = \mu_0$. But how small is small?

Standardizing the difference $\bar{y} - \mu_0$ by an estimate of $\text{sd}(\bar{Y}) = \sigma/\sqrt{n}$, namely the *standard error* of \bar{Y} , $\text{se}(\bar{Y}) = s/\sqrt{n}$ gives us a known distribution for the test statistic t^* *before we collect data*.

Reminder

If $\mu = \mu_0$ is true, then $t^* \sim t_{n-1}$.

Three types of test

Two sided: $H_0 : \mu = \mu_0$ versus $H_a : \mu \neq \mu_0$.

One sided, “<”: $H_0 : \mu = \mu_0$ versus $H_a : \mu < \mu_0$.

One sided, “>”: $H_0 : \mu = \mu_0$ versus $H_a : \mu > \mu_0$.

If the t^* we observe is highly unusual (relative to what we might see for a t_{n-1} distribution), we may reject H_0 and conclude H_a .

Let α be the significance level of the test, the maximum allowable probability of rejecting H_0 when H_0 is true.

Rejection rules

- Two sided: If $|t^*| > t_{n-1}(1 - \alpha/2)$ then reject H_0 , otherwise fail to reject H_0 .
- One sided, $H_a : \mu < \mu_0$. If $t^* < t_{n-1}(\alpha)$ then reject H_0 , otherwise fail to reject H_0 .
- One sided, $H_a : \mu > \mu_0$. If $t^* > t_{n-1}(1 - \alpha)$ then reject H_0 , otherwise fail to reject H_0 .

P-value approach

We can also measure the evidence against H_0 using a p-value, which is the probability of observing a test statistic value *as extreme or more extreme* than the test statistic *we did observe*, if H_0 were true.

A small p-value provides strong evidence against H_0 .


Rule: p-value $< \alpha \Rightarrow$ reject H_0 , otherwise accept H_0 .

p-values are computed according to the alternative hypothesis.

Let $T \sim t_{n-1}$; then

- Two sided: $2 \times \min\{P(T < t^*), P(T > t^*)\}$
- One sided, $H_a : \mu < \mu_0$: $p = P(T < t^*)$.
- One sided, $H_a : \mu > \mu_0$: $p = P(T > t^*)$.

Example



Example: We wish to test whether the true mean high temperature is greater than 75° using $\alpha = 0.01$:

$$H_0 : \mu = 75 \text{ versus } H_a : \mu > 75.$$

$$t^* = \frac{77.667 - 75}{8.872/\sqrt{30}} = 1.646 < t_{29}(0.99) = 2.462.$$

What do we conclude? Note that $p = 0.05525 > 0.01$.

- Consider that we do not reject H_0 if

$$\left| \frac{\bar{X} - \mu_0}{s/\sqrt{n}} \right| \leq t_{1-\alpha/2, n-1}$$

implying

$$|\bar{X} - \mu_0| \leq t_{1-\alpha/2, n-1} s/\sqrt{n}$$

implying

$$\bar{X} - t_{1-\alpha/2, n-1} s/\sqrt{n} < \mu_0 < \bar{X} + t_{1-\alpha/2, n-1} s/\sqrt{n}$$

Connection between CI and two-sided test

An α -level two-sided test rejects $H_0 : \mu = \mu_0$ if and only if μ_0 falls outside the $(1 - \alpha)100\%$ CI about μ .

Example (continued): Recall that the 90% CI for New York's high temperature is (74.91,80.42) degrees.

- At $\alpha = 0.10$, would we reject $H_0 : \mu = 73$ and conclude $H_a : \mu \neq 73$?
- At $\alpha = 0.10$, would we reject $H_0 : \mu = 80$ and conclude $H_a : \mu \neq 80$?
- At $\alpha = 0.05$, would we reject $H_0 : \mu = 80$ and conclude $H_a : \mu \neq 80$?

R Code

```
> summer.temps <- c(59,81,85,75,81,77,92,83,92,76,74,73,57,68,80,69,  
+ 83,78,87,88,93,71,73,76,81,82,73,64,80,79)  
> mean(summer.temps)  
[1] 77.66667  
> sd(summer.temps)  
[1] 8.872015  
> t.test(summer.temps, mu=75, conf.level=0.9)  
One Sample t-test  
  
data:  summer.temps  
t = 1.6463, df = 29, p-value = 0.1105  
alternative hypothesis: true mean is not equal to 75  
90 percent confidence interval:  
 74.91442 80.41891  
sample estimates:  
mean of x  
 77.66667
```

Example: $n = 7$ pairs of mice were injected with a cancer cell. Mice within each pair came from the same litter and were therefore biologically similar. For each pair, one mouse was given an experimental drug and the other mouse was untreated. After a time, the mice were sacrificed (killed) and the tumors weighed.

When we have two paired samples (when each observation in one sample can be naturally paired with an observation in the other sample), we can use one-sample methods to obtain inference on the **mean difference**.

One-sample inference on differences

Let (Y_{1j}, Y_{2j}) be the pair of tumor weights from control and treatment mice within litter j , $j = 1, \dots, 7$.

The difference of tumor weights in control versus treatment within each litter is

$$D_j = Y_{1j} - Y_{2j}.$$

If the differences follow a normal distribution, then we have the model

$$D_j = \mu_D + e_j, \quad j = 1, \dots, n, \quad \text{where } e_1, \dots, e_n \stackrel{iid}{\sim} N(0, \sigma^2).$$


Note that μ_D is the mean difference.

To test whether the control results in a higher mean tumor weight, form

$$H_0 : \mu_D = 0 \text{ versus } H_a : \mu_D > 0.$$

```
ods graphics on;
data mice plots;
input control treatment @@;
datalines;
1.321 0.841 1.423 0.932 2.682 2.011 0.934 0.762 1.230 0.991 1.670
1.120 3.201 2.312
;
proc ttest h0=0 alpha=0.01 sides=u;
paired control*treatment;
run;
ods graphics;
```

Paired example, continued



For this test, the p-value is 0.0008. At $\alpha = 0.05$, we reject H_0 and conclude that the true mean difference of tumor weights is greater than 0.

Restated: the treatment produces a significantly lower mean tumor weight.

A 95% CI for the true mean difference μ_D is (0.27, 0.73). With 95% confidence, the mean tumor weight for untreated mice is between 0.27 and 0.73 grams higher than for treated mice.

Section A.7 Two independent samples

Assume we have two *independent* (not paired) samples from two normal populations. Label them 1 and 2. The model is

$$Y_{ij} = \mu_i + \epsilon_{ij}, \text{ where } i = 1, 2 \text{ and } j = 1, \dots, n_i.$$

The “within sample heterogeneity” follows

$$\epsilon_{ij} \stackrel{iid}{\sim} N(0, \sigma^2).$$

Section A.7 Two independent samples

Assume we have two *independent* (not paired) samples from two normal populations. Label them 1 and 2. The model is

$$Y_{ij} = \mu_i + \epsilon_{ij}, \text{ where } i = 1, 2 \text{ and } j = 1, \dots, n_i.$$

The “within sample heterogeneity” follows

$$\epsilon_{ij} \stackrel{iid}{\sim} N(0, \sigma^2).$$

- Both populations have the same variance σ^2 .

Section A.7 Two independent samples

Assume we have two *independent* (not paired) samples from two normal populations. Label them 1 and 2. The model is

$$Y_{ij} = \mu_i + \epsilon_{ij}, \text{ where } i = 1, 2 \text{ and } j = 1, \dots, n_i.$$

The “within sample heterogeneity” follows

$$\epsilon_{ij} \stackrel{iid}{\sim} N(0, \sigma^2).$$

- Both populations have the same variance σ^2 .
- The two sample sizes (n_1 and n_2) may be different.

Pooled approach

An estimator of the variance σ^2 is the “pooled sample variance”

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}.$$

Then

$$t = \frac{(\bar{Y}_1 - \bar{Y}_2) - (\mu_1 - \mu_2)}{\sqrt{s_p^2 \left[\frac{1}{n_1} + \frac{1}{n_2} \right]}} \sim t_{n_1+n_2-2}.$$

We are interested in the mean difference $\mu_1 - \mu_2$, i.e. the difference in the population means.

A $(1 - \alpha)100\%$ CI for $\mu_1 - \mu_2$ is

$$(\bar{Y}_1 - \bar{Y}_2) \pm t_{n_1+n_2-2}(1 - \alpha/2) \sqrt{s_p^2 \left[\frac{1}{n_1} + \frac{1}{n_2} \right]}.$$

Pooled approach: Hypothesis test

Often we wish to test whether the two populations have the same mean, i.e. $H_0 : \mu_1 = \mu_2$. Of course, this implies $H_0 : \mu_1 - \mu_2 = 0$. The test statistic is

$$t^* = \frac{\bar{Y}_1 - \bar{Y}_2}{\sqrt{s_p^2 \left[\frac{1}{n_1} + \frac{1}{n_2} \right]}}$$

and is distributed $t_{n_1+n_2-2}$ under H_0 . Let $T \sim t_{n_1+n_2-2}$. The tests are carried out via:

H_a	Rejection rule	p-value
$\mu_1 \neq \mu_2$	$ t^* > t_{n_1+n_2-2}(1 - \alpha/2)$	$P(T > t^*)$
$\mu_1 < \mu_2$	$t^* < -t_{n_1+n_2-2}(1 - \alpha)$	$P(T < t^*)$
$\mu_1 > \mu_2$	$t^* > t_{n_1+n_2-2}(1 - \alpha)$	$P(T > t^*)$

Unequal variances: Satterthwaite approximation

What if it is not reasonable to assume the populations have the same variance—i.e., $\sigma_1^2 \neq \sigma_2^2$? The model is

$$Y_{ij} = \mu_i + \epsilon_{ij}, \quad \epsilon_{ij} \stackrel{\text{ind.}}{\sim} N(0, \sigma_i^2)$$

where $i = 1, 2$ denotes the population and $j = 1, \dots, n_i$ the measurement within the population.

Use s_1^2 and s_2^2 to estimate σ_1^2 and σ_2^2 . Define the test statistic

$$t^* = \frac{\bar{Y}_1 - \bar{Y}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}.$$

Unequal variances: Satterthwaite approximation

Under the null $H_0 : \mu_1 = \mu_2$, this test statistic is *approximately* distributed $t^* \sim t_{df}$ where

$$df = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{s_1^4}{n_1^2(n_1-1)} + \frac{s_2^4}{n_2^2(n_2-1)}}.$$

Note that $df = n_1 + n_2 - 2$ when $n_1 = n_2$ and $s_1 = s_2$.

Satterthwaite and pooled variance methods typically give similar results when $s_1 \approx s_2$.

Testing $H_0 : \sigma_1 = \sigma_2$

- We can formally test $H_0 : \sigma_1 = \sigma_2$ using Bartlett's F-test or Levene's test, but in practice graphical methods such as box plots are often used.
- The "var.test" function in R


$$F^* = \frac{\max\{s_1^2, s_2^2\}}{\min\{s_1^2, s_2^2\}}$$

- This test assumes normal data and is sensitive to this assumption.

Two sample T test example

Example: Data were collected on pollution around a chemical plant (Rao, p. 137). Two independent samples of river water were taken, one upstream and one downstream. Pollution level was measured in ppm. Do the mean pollution levels differ at $\alpha = 0.05$?

```
upstream <- c(24.5, 29.7, 20.4, 28.5, 25.3, 21.8, 20.2, 21.0, 21.9, 22.2)
downstream <- c(32.8, 30.4, 32.3, 26.4, 27.8, 26.9, 29.0, 31.5,
31.2, 26.7, 25.6, 25.1, 32.8, 34.3, 35.4)
boxplot(upstream, downstream)
# Do the spreads seem equal across groups?
```



Pollution Example (Continued)

```
> # Assuming equal variances:  
> t.test(upstream, downstream, var.equal=TRUE)
```

Two Sample t-test

```
data: upstream and downstream  
t = -4.6433, df = 23, p-value = 0.0001132  
alternative hypothesis: true difference in means is not equal to 0  
95 percent confidence interval:  
 -9.150113 -3.509887  
sample estimates:  
mean of x mean of y  
 23.55      29.88  
> # Not assuming equal variances:  
> t.test(upstream, downstream)
```

Welch Two Sample t-test

```
data: upstream and downstream  
t = -4.6339, df = 19.31, p-value = 0.0001742  
alternative hypothesis: true difference in means is not equal to 0  
95 percent confidence interval:  
 -9.186009 -3.473991  
sample estimates:  
mean of x mean of y  
 23.55      29.88  
> # A 99% CI for the difference in means:  
> t.test(upstream, downstream, var.equal=TRUE, conf.level=0.99)$conf.int  
[1] -10.157121 -2.502879
```


T procedure assumptions

Note: Recall our t-procedures require that the data come from normal population(s).

Fortunately, the t procedures are **robust**: they work approximately correctly if the population distribution is “close” to normal.

Also, if our sample sizes are large, we can use the t procedures (or simply normal-based procedures) even if our data are not normal because of the central limit theorem.

If the sample size is small, we should perform some check of normality to ensure t tests and CIs are okay.

Question: Are there any other model assumptions that can or should be checked? For example, what if pollution measurements were taken on consecutive days?

Boxplots for checking normality

- To use t tests and CIs in small samples, approximate normality should be checked.
- We can check with a histogram or boxplot: verify distribution is unimodal and *approximately* symmetric.
- **Note:** For normal data, the probability of seeing an outlier on a R using defaults is 0.0070. For a sample size $n_j = 150$ from normal data, we expect to see $0.0070 \times 150 \approx 1$ outlier. Certainly for small sample sizes, we expect to see none.

Q-Q plot for checking normality

- A more precise plot: normal Q-Q plot. Idea: the human eye is very good at detecting deviations from linearity.
- Plot ordered data $\{y_{(i)}\}$ against normal quantiles $z_i = \Phi^{-1}\{i/(n+1)\}$ for $i = 1, \dots, n$.
- Idea: $z_i \approx E(Z_{(i)})$, the expected order statistic under standard normal assumption.
- A plot of $y_{(i)}$ versus z_i should be reasonably straight *if data are normal*.
- However, in small sample sizes there is a lot of variability in the plots even with perfectly normal data...

What to Use

The t-test relies on a normality assumption. When sample size is small, consider:

- Wilcoxon Rank Sum Test
- Permutation Test

→ The crucial assumption is independence between observations.