

Semiparametric Estimation in the Secondary Analysis of Case-Control Studies

Yanyuan Ma and Raymond J. Carroll

Department of Statistics, University of South Carolina, Columbia, SC 29208

Department of Statistics, Texas A&M University, College Station, TX 77843

yanyuan.ma@stat.sc.edu and carroll@stat.tamu.edu

Abstract

We study the regression relationship among covariates in case-control data, an area known as the secondary analysis of case-control studies. The context is such that only the form of the regression mean is specified, so that we allow an arbitrary regression error distribution, which can depend on the covariates and thus can be heteroscedastic. Under mild regularity conditions we establish the theoretical identifiability of such models. Previous work in this context has either (a) specified a fully parametric distribution for the regression errors, (b) specified a homoscedastic distribution for the regression errors, (c) has specified the rate of disease in the population (we refer this as true population), or (d) has made a rare disease approximation. We construct a class of semiparametric estimation procedures that rely on none of these. The estimators differ from the usual semiparametric ones in that they draw conclusions about the true population, while technically operating in a hypothetical superpopulation. We also construct estimators with a unique feature, in that they are robust against the misspecification of the regression error distribution in terms of variance structure, while all other nonparametric effects are estimated despite of the biased samples. We establish the asymptotic properties of the estimators and illustrate their finite sample performance through simulation studies, as well as through an empirical example on the relation between red meat consumption and heterocyclic amines. Our analysis verified the positive relationship between red meat consumption and two forms of HCA, indicating that increased red meat consumption leads to increased levels of MeIQa and PhIP, both being risk factors for colorectal cancer. Computer software as well as data to illustrate the methodology are available at http://www.stat.tamu.edu/~carroll/matlab_programs/software.php.

Some Key Words: Biased samples; Case-control study; Heteroscedastic regression; Secondary analysis; Semiparametric estimation.

Short title: Secondary Analysis of Case-Control Studies

1 Introduction

Population-based case-control designs, hereafter called case-control designs, are popularly used for studying risk factors for rare diseases, such as cancers. The idealized set up of such designs is as follows. At a given time, there is an underlying base population, which we refer to as the *true population* throughout the paper. Within the true population, there are two subpopulations, those with the disease, called cases, and those without the disease, called controls. Separately, a random sample is taken from the case subpopulation, and a random sample is taken from the control subpopulation. Data on various covariates are then collected in a retrospective fashion, so that they reflect history prior to the disease. Nested case-control studies and case-cohort or case-base studies are variations of the retrospective case-control design.

The primary purpose of case-control designs is to understand the relation between disease occurrence and the covariates. The *secondary analysis* of such case-control data (Jiang et al., 2006; Lin and Zeng, 2009; Li et al., 2010; Wei et al., 2013; He et al., 2012) is based on the realization that the data further provide information about the relationship among the covariates. The relation between covariates are often of interest as well, as they can reveal associations between various covariates such as gene-environment, gene-gene and environment-environment associations. These analyses become especially important when, as is the case of retrospective sampling, a random sample from the true population is not available; see the secondary analysis literature mentioned above for more examples. If we seek to understand the regression relationship between covariates Y and \mathbf{X} in the true population, we generally cannot use the case-control data set as if it were a random sample from the true population. Indeed, unless disease is independent of Y given \mathbf{X} , the regression of Y on \mathbf{X} based on the case-control sample will lead to a relationship different from that in the true population.

To see this numerically, we first define our notation. There are N_0 cases and N_1 controls, with $N = N_0 + N_1$. Suppose that $N_0 = N_1 = 500$, and that disease status D is related to covariates (Y, \mathbf{X}) in the true population through the linear logistic model

$$\text{pr}(D = d | \mathbf{X} = x, Y = y) = f_{D|X,Y}^{\text{true}}(d, \mathbf{x}, y) = H(d, \mathbf{x}, y, \boldsymbol{\alpha}) = \frac{\exp\{d(\alpha_c + \mathbf{x}^T \boldsymbol{\alpha}_1 + y\alpha_2)\}}{1 + \exp(\alpha_c + \mathbf{x}^T \boldsymbol{\alpha}_1 + y\alpha_2)}, \quad (1)$$

where for this illustration, $\boldsymbol{\alpha} = (\alpha_c, \boldsymbol{\alpha}_1, \alpha_2) = (-5.5, 1.0, 0.5)$. Suppose further that the

regression relationship in the true population is that $Y = \beta_c + X\beta + \epsilon$, with $\beta_c = 0$, $\beta = 1$ and $\epsilon \sim \text{Normal}(0, 1)$. In addition, in the true population, $X \sim \text{Uniform}(0, 1)$. In this setup, suppose the disease is rare, with $\text{pr}(D = 1) \approx 0.01$. Thus, while controls are 99% of the true population, they are only 50% of the case-control study. To understand the bias induced by ignoring the case-control sampling scheme, we generated 3,000 case-control studies with intercept $\beta_c = 0$ and slope $\beta = 1$, and computed the intercept and slope estimates using all the data. Simply regressing Y on X and ignoring the case-control sampling scheme, the mean estimated intercept and slope across the 3,000 simulated data sets were 0.150 and 1.174, respectively, reflecting considerable bias, which leads to a coverage rate of only 67% for a nominal 95% confidence interval. Figure 1 shows the attained regression function compared to the true regression function. Using the method that we develop in this paper, our method yields the average intercept and slope estimates of 0.0024 and 1.0035, thus eliminating the bias caused by ignoring the case-control sampling scheme.

The bias in the secondary analysis is in stark contrast to what happens in the primary analysis, where estimating (α_1, α_2) is of interest. It is well known that α_1 and α_2 can be estimated consistently via ordinary logistic regression of D on (Y, \mathbf{X}) by treating the case-control sample as if it were a random sample of the true population (Prentice and Pyke, 1979).

Our goal is to estimate the regression of Y on \mathbf{X} in the true population, using case-control data, where for a function $m(\cdot)$ known up to a parameter β ,

$$Y = m(\mathbf{X}, \beta) + \epsilon, \tag{2}$$

where we make only the assumption that $E(\epsilon|\mathbf{X}) = 0$. Two solutions to estimating β have been proposed in the literature. (Lin and Zeng, 2009) and, obliquely, (Chen et al., 2008) proposed to assume a particular fully parametric distribution for ϵ and then perform a semi-parametric efficient analysis, where the distribution of \mathbf{X} is nonparametric. There is excellent software for this problem in the case that $\epsilon = \text{Normal}(0, \sigma^2)$, i.e., homoscedastic and normally distributed (<http://www.bios.unc.edu/~lin/software/SPREG/>). To implement this software, however, one must either specify the disease rate $\text{pr}(D = 1)$ in the true population or one must make a “rare-disease” assumption, which is implemented by assuming

$\text{pr}(D = 1) < 0.01$. When the disease rate is known, reweighting the observations also corrects the biases (Scott and Wild, 2002). Wei, et al. (2012) dispense with the normality assumption, but still assume a homoscedastic distribution for ϵ independent of \mathbf{X} and make a rare disease approximation.

In practice, the disease rate in the population being sampled is not known. In addition, it might not be rare. As an example, in Section 6, we use data from a case-control study of colorectal adenoma, a precursor to colorectal cancer, relating measures of heterocyclic amines to red meat consumption. While colorectal cancer is rare, colorectal adenomas are not, being on the order of 7% or more depending on the population being sampled (Yamaji et al., 2004; Corley et al., 2014). In this data set, one of the regressions is also heavily heteroscedastic. We will demonstrate that both approaches mentioned above have problems when some of the assumptions, such as the rare disease assumption, the known disease rate assumption and the known error distribution assumption, are violated (Tables 1-6).

In order to relax such assumptions, novel methods are needed. In this paper, we do not assume any distributional form for ϵ or $\epsilon \mid \mathbf{X}$, we do not assume that the regression is homoscedastic, we do not require the disease rate to be known and we do not make a rare disease approximation. We do this by adopting the concept of a superpopulation (Ma, 2010): a similar idea is called an alternative characterization of the case-control study by Chen et al. (2009).

The main idea behind a superpopulation is to enable us to view the case-control sample as a sample of independent and identically distributed (iid) observations from the superpopulation. Conceptually, superpopulation is simply a proportional expansion of the case-control sample to infinity. Why a superpopulation constructed through such expansion achieves the purpose of viewing the case-control sample as an iid sample is studied carefully Ma (2010). The ability of viewing the case-control sample as a random sample permits us to use classical semiparametric approaches (Bickel et al., 1993; Tsiatis, 2006), regardless if the disease rate in the real population is rare or not, or is known or not.

We derive a class of semiparametric estimators and identify the efficient member. We further construct a member of the family that is relatively simple to compute, and illustrate how to construct the efficient estimator, applicable to both rare and common diseases. The

derivation of semiparametric estimators in this context is challenging because the calculations must use quantities defined in the unknown true population to perform analysis in the superpopulation, since the models under the true population and the superpopulation share common parameters. In addition, as established in Ma (2010), the resulting semiparametric estimators further retain asymptotic consistency, a root- n convergence rate, asymptotic normality and semiparametric efficiency with respect to the true population as well. For example, our efficient estimator has the usual property that its asymptotic variance cannot be further reduced by any other device or by taking into account the case-control sampling structure.

The rest of the paper is organized as follows. Under conditions, we first establish the technical identifiability of our problem in Section 2. In Section 3, we formulate the problem into a classic semiparametric one by using the superpopulation notion and carry out analytic calculations to prepare for the estimation procedure. In Section 4, we describe details of implementation and the asymptotic theory. Simulation studies are performed in Section 5 to illustrate the finite sample performance of the procedure, showing that our method is robust, efficient and maintains nominal coverage for confidence intervals. An empirical analysis is provided in Section 6. Section 7 contains a short discussion. Technical details are given in an Appendix, as well as in the **Supplementary Material**. Computer code and data to illustrate our method are available http://www.stat.tamu.edu/~carroll/matlab_programs/software.php.

2 The Superpopulation Model Framework

The primary disease model is the linear logistic model (1), with $\boldsymbol{\alpha} = (\alpha_c, \boldsymbol{\alpha}_1^T, \alpha_2)^T$. Here and throughout the text, we use superscript “true” to represent quantities or operations related to the underlying true population, and also to distinguish it from a superpopulation that will be formally introduced later. In addition, in this underlying true population, Y is believed to be related to \mathbf{X} through (2), which we rewrite as the regression model

$$f_{Y|X}^{\text{true}}(\mathbf{x}, y) = \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\}, \quad (3)$$

where $m(\cdot)$ is the regression mean function known up to the parameter $\boldsymbol{\beta}$ and η_2 is an unknown probability density function that has mean zero given \mathbf{X} . Defining $\epsilon = Y - m(\mathbf{X}, \boldsymbol{\beta})$, then $E(\epsilon | \mathbf{X}) = 0$. The distribution of ϵ , whether conditional on \mathbf{X} or marginally, is left unspecified. In particular, heteroscedasticity is allowed. Making the identification $\eta_2(\epsilon, \mathbf{X}) = \eta_2\{Y - m(\mathbf{X}, \boldsymbol{\beta}), \mathbf{X}\}$, this means that $\eta_2 \geq 0$ satisfies $\int \epsilon \eta_2(\epsilon, \mathbf{x}) d\mu(\epsilon) = 0$ and $\int \eta_2(\epsilon, \mathbf{x}) d\mu(\epsilon) = 1$, but its form is unknown. Here and throughout the text, we use $\mu(\cdot)$ to denote a Lebesgue measure for a continuous random variable and a counting measure for a discrete random variable. The distribution of the covariate \mathbf{X} in the underlying true population is also unspecified, and its density or mass function is $f_{\mathbf{X}}^{\text{true}}(\mathbf{x}) = \eta_1(\mathbf{x})$, where $\eta_1 \geq 0$ satisfies $\int \eta_1(\mathbf{x}) d\mu(\mathbf{x}) = 1$.

The superpopulation framework of Ma (2010) is that one can think of the case-control sample as a random sample from an imaginary infinite superpopulation, in which the disease to non-disease ratio is N_1/N_0 . Let $N_d = N_0$ when $d = 0$ and $N_d = N_1$ when $D = 1$. Define the true probability that $D = d$ as $p_D^{\text{true}}(d, \boldsymbol{\alpha}, \boldsymbol{\beta}, \eta_1, \eta_2) = \int \eta_1(\mathbf{x}) \eta_2(\epsilon, \mathbf{x}) H(d, \mathbf{x}, y, \boldsymbol{\alpha}) d\mu(\mathbf{x}) d\mu(y)$. The density of (D, Y, \mathbf{X}) in the superpopulation is defined as

$$f_{X,Y,D}(\mathbf{x}, y, d) = \frac{N_d \eta_1(\mathbf{x}) \eta_2(\epsilon, \mathbf{x}) H(d, \mathbf{x}, y, \boldsymbol{\alpha})}{N p_D^{\text{true}}(d, \boldsymbol{\alpha}, \boldsymbol{\beta}, \eta_1, \eta_2)}. \quad (4)$$

Although $\boldsymbol{\beta}$ appears in ϵ , for notational brevity, we do not explicitly write $\epsilon(\boldsymbol{\beta})$. In the secondary analysis framework, the main interest is $\boldsymbol{\beta}$. However we formally treat $\boldsymbol{\theta} = (\boldsymbol{\alpha}^T, \boldsymbol{\beta}^T)^T$ as the parameter of interest. We treat $\eta_1(\cdot)$ and $\eta_2(\cdot, \cdot)$ as the infinite dimensional nuisance parameters, thus bypassing the need to estimate them.

Remark 1. When no assumptions are made about the relationship between Y and X in the true population, the logistic intercept α_c is not identified (Prentice and Pyke, 1979), and neither is the regression of Y on X . Thus, if consistency of estimation is desired, truly nonparametric regression in a case-control study of our type is not possible. We believe that the key to identification lies in placing a restriction on the joint distribution of (Y, X) in the base population. For example, Chatterjee and Carroll (2005) show that if Y and X are independent, then α_c is generally identified, and they show this explicitly when one of the two is discrete. In our case, the restriction is a parametric model for $E(Y|X)$. It

is a reasonable conjecture that such a restriction is enough for the identifiability of α_c , a conjecture that we confirm next.

2.1 Identifiability

We first establish identifiability of the parameters $\boldsymbol{\alpha}, \boldsymbol{\beta}$ in the superpopulation. For greater generality, we consider the slightly more flexible model $H(d, \mathbf{x}, y) = \exp[d\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}]/[1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}]$, where $u(0, 0, \boldsymbol{\alpha}_1, \alpha_2) = 0$ for all $\boldsymbol{\alpha}_1, \alpha_2$. Obviously, this model contains the original linear logistic model we are studying. We assume that there is no $(\boldsymbol{\alpha}_1^T, \alpha_2)^T \neq (\tilde{\boldsymbol{\alpha}}_1^T, \tilde{\alpha}_2)^T$ such that for all (\mathbf{x}, y) , $u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2) = u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)$. These are natural minimal conditions that are usually satisfied automatically as long as the parameterizations of u and m are not redundant. We also assume the following two conditions.

Assumption 1. Assume that the second moment of ϵ is bounded marginally and η_2 is a bounded function, i.e., $E(\epsilon^2) < \infty$ and $\sup_{\mathbf{x}, \epsilon} \eta_2(\epsilon, \mathbf{x}) < \infty$. For any fixed parameters $\boldsymbol{\alpha}_1, \alpha_2, \boldsymbol{\beta}$, and any $\delta > 0$, there exists a constant vector \mathbf{c}_1 , a constant $c_2 \in [0, 1]$ and a region \mathcal{D} with complement \mathcal{D}^c such that when $\mathbf{x} \rightarrow \mathbf{c}_1$,

$$\sup_{\epsilon \in \mathcal{D}^c} \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} |(1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}])^{-1} - c_2| = 0,$$

and $\lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \text{pr}(\epsilon \in \mathcal{D} \mid \mathbf{X} = \mathbf{x}) < \delta$. In addition, for any element $e \in \mathcal{D}$, $|e| \geq 1$. Typically we expect $\mathcal{D}^c = [-K, K]$ for some large K , $\mathbf{c}_1 = \infty$ or $-\infty$ or contains $\pm\infty$ as components, and $c_2 = 0$ or 1 , although this is not required.

Assumption 2. $c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) = \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \{m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) - m(\mathbf{x}, \boldsymbol{\beta})\} \neq 0$ for $\tilde{\boldsymbol{\beta}} \neq \boldsymbol{\beta}$.

Remark 2. Assume that $\text{pr}(|\epsilon| > K \mid \mathbf{X} = \mathbf{x}) \rightarrow 0$ as $K \rightarrow \infty$ uniformly in \mathbf{x} . We can easily verify that when both m and u are linear functions, where we write $m(\mathbf{x}, \boldsymbol{\beta}) = \mathbf{x}^T \boldsymbol{\beta}_1 + \beta_c$, both assumptions are satisfied except when $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2 = \mathbf{0}$. When this happens, $u\{m(\mathbf{x}, \boldsymbol{\beta}), \boldsymbol{\alpha}_1, \alpha_2\}$ degenerates to a constant, and we can verify that although $\boldsymbol{\beta}_1$ is still identifiable, β_c and α_c are no longer identifiable, see the **Supplementary Material** for details of verification of both the identifiability and the non-identifiability verification.

We state the identifiability result in Proposition 1 and provide the proof in Appendix A.1.

Proposition 1. *Make Assumptions 1-2. Also assume that there are constants (C_1, C_2) such that $0 < C_1 < N_0/N_1 < C_2 < \infty$. Then the parameters $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are identifiable.*

Remark 3. Identifiability under some specific situations has been considered in the literature. For example, Chatterjee and Carroll (2005), Chatterjee et al. (2006) and Chen et al. (2009) considered the case that \mathbf{X} and Y are independent, while Chen et al. (2008) and Lin and Zeng (2009) explicitly studied the identifiability issue when the disease rate model is linear logistic and the secondary model is fully parametric. The model we consider here is more general, in that only a mean function is assumed for the secondary model. These authors all note that while in practice, it may be difficult to estimate α_c , estimation of the other parameters can still be performed effectively, see also Lobach et al. (2008).

3 Analytic Derivations

3.1 True and Conjectured Models

The major point of our article is that we only propose a model for $E(Y | \mathbf{X})$, denoted $m(\mathbf{X}, \boldsymbol{\beta})$, and we specifically want to avoid positing a model for the density function of the regression errors $\epsilon = Y - m(\mathbf{X}, \boldsymbol{\beta})$ conditional on \mathbf{X} . We will accomplish this by a two-step process. First, in Section 3.2, we will derive the semiparametric efficient estimating equation in the superpopulation for estimating $(\boldsymbol{\alpha}, \boldsymbol{\beta})$ when the density of Y given X in the true population is known. Recognizing that we do not want to make such an assumption, in Section 4, we will show how to modify the estimating equation so that it has mean zero asymptotically, even if the conjectured model for the regression errors is false, thus resulting in model-robust consistent estimation.

3.2 Analysis Under a True Model

As described in Section 3.1, here we will derive the form of the semiparametric efficient estimating equation when the conjectured model for the regression errors in (3) is true. Later in Section 4, we will modify the estimating function to make it model-robust.

Viewing the observations as randomly sampled from the superpopulation, we can perform a conventional semiparametric analysis. Of course, all the calculations need to be done with respect to the superpopulation, and all the probability statements need to be with respect to Lebesgue measure for continuous random variables and counting measure for discrete random variables in the superpopulation, and they will be if not otherwise pointed out. The functions (η_1, η_2, H) , which are probability density/mass functions in the true population, do not represent the corresponding probabilities density/mass functions in the superpopulation. They are merely functions that satisfy $\eta_1(\mathbf{x}) \geq 0, \int \eta_1(\mathbf{x})d\mu(\mathbf{x}) = 1, \eta_2(\epsilon, \mathbf{x}) \geq 0, \int \eta_2(\epsilon, \mathbf{x})d\mu(\epsilon) = 1, \int \epsilon\eta_2(\epsilon, \mathbf{x})d\mu(\epsilon) = 0, H(d, \mathbf{x}, y) \geq 0, H(0, \mathbf{x}, y) + H(1, \mathbf{x}, y) = 1$. In fact, we introduced these symbols to discourage the mistake of automatically viewing them as the corresponding density or mass functions in the superpopulation.

Using model (4), calculating the partial derivative of the loglikelihood with respect to $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$, it is easy to see that the score function has the form $\mathbf{S}_\theta(\mathbf{X}, Y, D, \boldsymbol{\theta}) = \mathbf{S}(\mathbf{X}, Y, D, \boldsymbol{\theta}) - E(\mathbf{S} | D)$, where $\boldsymbol{\theta} = (\boldsymbol{\alpha}^\top, \boldsymbol{\beta}^\top)^\top$, $\mathbf{S}_\theta = (\mathbf{S}_\alpha^\top, \mathbf{S}_\beta^\top)^\top$, and

$$\mathbf{S}(\mathbf{x}, y, d, \boldsymbol{\theta}) = \begin{Bmatrix} \partial \log\{H(d, \mathbf{x}, y, \boldsymbol{\alpha})\}/\partial \boldsymbol{\alpha} \\ \partial \log\{\eta_2(\epsilon, \mathbf{x})\}/\partial \boldsymbol{\beta} \end{Bmatrix}. \quad (5)$$

Explicitly,

$$\mathbf{S}_\alpha(\mathbf{X}, Y, D, \boldsymbol{\theta}) = \partial \log\{H(D, \mathbf{X}, Y, \boldsymbol{\alpha})\}/\partial \boldsymbol{\alpha} - E[\partial \log\{H(D, \mathbf{X}, Y, \boldsymbol{\alpha})\}/\partial \boldsymbol{\alpha} | D];$$

$$\mathbf{S}_\beta(\mathbf{X}, Y, D, \boldsymbol{\theta}) = \partial \log\{\eta_2(\epsilon, \mathbf{X})\}/\partial \boldsymbol{\beta} - E[\partial \log\{\eta_2(\epsilon, \mathbf{X})\}/\partial \boldsymbol{\beta} | D].$$

In Appendix A.2, we further derive the nuisance tangent space Λ and its orthogonal complement space Λ^\perp as

$$\begin{aligned} \Lambda &= \{\mathbf{g}(\epsilon, \mathbf{X}) - E(\mathbf{g} | D) : E_{\text{true}}(\mathbf{g}) = E_{\text{true}}(\epsilon \mathbf{g} | \mathbf{X}) = \mathbf{0} \text{ a.s.}\}; \\ \Lambda^\perp &= [\mathbf{h}(D, \epsilon, \mathbf{X}) : E(\mathbf{h}) = \mathbf{0}, E\{\mathbf{h} - E(\mathbf{h} | D) | \epsilon, \mathbf{X}\} \\ &\quad \times \sum_d (N_d/N)H(d, \mathbf{X}, y)/p_D^{\text{true}}(d) = \epsilon \mathbf{a}(\mathbf{X}) \text{ a.s.}], \end{aligned}$$

where $\mathbf{g}(\epsilon, \mathbf{x})$ and $\mathbf{h}(D, \epsilon, \mathbf{x})$ are arbitrary functions that satisfy their respective constraints described above, $\mathbf{a}(\mathbf{x})$ is an arbitrary function of \mathbf{x} , and a.s. stands for almost surely with respect to the true superpopulation distribution.

Having obtained both the score function and the two spaces Λ and Λ^\perp , conceptually, we only need to project the score function onto Λ^\perp to obtain the efficient score \mathbf{S}_{eff} . Doing this is, however, extraordinarily technical, and hence we defer the details to the **Supplementary Material**. Here we merely state the result in Proposition 2, which requires a series of definitions, as follows.

$$\begin{aligned}
& \text{Define } \pi_0 = p_D^{\text{true}}(0) = \int \eta_1(\mathbf{x})\eta_2(\epsilon, \mathbf{x})H(0, \mathbf{x}, y)d\mu(\mathbf{x})d\mu(y); \\
& \pi_1 = p_D^{\text{true}}(1) = \int \eta_1(\mathbf{x})\eta_2(\epsilon, \mathbf{x})H(1, \mathbf{x}, y)d\mu(\mathbf{x})d\mu(y); \\
& b_0 = E\{f_{D|X,Y}(1, \mathbf{X}, Y) \mid D = 0\}; b_1 = E\{f_{D|X,Y}(0, \mathbf{X}, Y) \mid D = 1\}; \\
& \mathbf{c}_0 = E(\mathbf{S} \mid D = 0) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 0\}; \\
& \mathbf{c}_1 = E(\mathbf{S} \mid D = 1) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 1\}; \\
& \kappa(\mathbf{x}, y) = [\sum_{d=0}^1 \{N_d H(d, \mathbf{x}, y)\} / (N\pi_d)]^{-1}; t_1(\mathbf{X}) = [E_{\text{true}} \{\epsilon^2 \kappa(\mathbf{X}, Y) \mid \mathbf{X}\}]^{-1}; \quad (6) \\
& \mathbf{t}_2(\mathbf{X}) = E_{\text{true}} \{\epsilon E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid \mathbf{X}\} - (\mathbf{c}_0/b_0)E_{\text{true}} \{\epsilon f_{D|X,Y}(0, \mathbf{X}, Y) \mid \mathbf{X}\}; \\
& t_3(\mathbf{X}) = -b_0^{-1}E_{\text{true}} \{\epsilon f_{D|X,Y}(0, \mathbf{X}, Y) \mid \mathbf{X}\}; \mathbf{a}(\mathbf{x}) = t_1(\mathbf{x})\{\mathbf{t}_2(\mathbf{x}) + t_3(\mathbf{x})\mathbf{u}_0\}; \\
& \mathbf{u}_0 = (1 - E[\epsilon t_1(\mathbf{X})t_3(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0])^{-1} E[\epsilon t_1(\mathbf{X})\mathbf{t}_2(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0]; \\
& \mathbf{u}_1 = -(N_0/N_1)\mathbf{u}_0; \mathbf{v}_0 = (\pi_1/b_0)(\mathbf{u}_0 + \mathbf{c}_0); \mathbf{v}_1 = -(\pi_0/b_0)(\mathbf{u}_0 + \mathbf{c}_0); \\
& \mathbf{g}(\epsilon, \mathbf{x}) = E(\mathbf{S} \mid \epsilon, \mathbf{X} = \mathbf{x}) - \epsilon \mathbf{a}(\mathbf{x})\kappa(\mathbf{x}, y) - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{x}, y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{x}, y).
\end{aligned}$$

Proposition 2. Make the definitions (6). In the superpopulation, the semiparametric efficient score function is $\mathbf{S}(\mathbf{X}_i, Y_i, D_i) - \mathbf{g}\{Y_i - m(\mathbf{X}_i, \boldsymbol{\beta}), \mathbf{X}_i\} - (N_0/N)\mathbf{v}_0 - (N_1/N)\mathbf{v}_1$. The semiparametric efficient estimator is obtained by solving

$$\sum_{i=1}^N [\mathbf{S}(\mathbf{X}_i, Y_i, D_i) - \mathbf{g}\{Y_i - m(\mathbf{X}_i, \boldsymbol{\beta}), \mathbf{X}_i\} - (1 - D_i)\mathbf{v}_0 - D_i\mathbf{v}_1] = \mathbf{0}. \quad (7)$$

We emphasize here that the estimator in Proposition 2 is not only efficient with respect to the superpopulation, it is also efficient with respect to the true population. This is a direct consequence of the general result that if an estimator is efficient with respect to the superpopulation, it is also efficient with respect to the true population. A careful justification of this claim is given in Ma (2010). Logically, this result can be understood because if we could find a more efficient estimator with respect to the true population, this estimator would also be more efficient with respect to the superpopulation, which causes a contradiction.

Intuitively, the special sampling strategy is in fact already absorbed into the formulation when we construct the superpopulation, hence no information has been lost during the conversion between populations.

4 Estimator Construction

4.1 Basic Calculations

The estimating equation (7) derived in Proposition 2 is not useful however, because it involves various calculations that rely on the unknown η_1 and η_2 , which were assumed to be correctly conjectured in Section 3. If either are misspecified, the corresponding calculation will lead to inconsistent estimation of $\boldsymbol{\theta}$. The purpose of this section is to define estimators that are consistent for estimating $\boldsymbol{\theta}$ based upon a posited score function, which we denote by \mathbf{S}^* . As it turns out, if the posited score function is correct, then in addition to being consistent, the estimator of $\boldsymbol{\theta}$ has the additional property of being efficient. If the posited score function is incorrect, then the estimator of $\boldsymbol{\theta}$ is still consistent. So our method can be thought of as a locally efficient estimator.

A careful inspection of the estimation procedure given in Proposition 2 and the definition of the related quantities suggests that the critical points lie in obtaining π_0 and π_1 , in calculating $E(\mathbf{h} \mid \epsilon, \mathbf{X})$ and $E(\mathbf{h} \mid D)$ for any function $\mathbf{h}(D, \mathbf{X}, Y)$, and in calculating $E_{\text{true}}(\mathbf{h} \mid \mathbf{X})$ for any function $\mathbf{h}(D, \mathbf{X}, Y)$.

Our algorithm is detailed as Algorithm 1, and is based upon the following considerations.

- First, we have that

$$\begin{aligned} N_d &= Np_D(d) = N \int f_{\mathbf{X},Y}(\mathbf{x}, y) f_{D|X,Y}(d, \mathbf{x}, y) d\mu(\mathbf{x}) d\mu(y) \\ &= N \int f_{\mathbf{X},Y}(\mathbf{x}, y) (N_d H / N\pi_d) \left\{ \sum_d (N_d H) / (N\pi_d) \right\}^{-1} d\mu(\mathbf{x}) d\mu(y). \end{aligned}$$

If we estimate the last term by $\sum_{i=1}^N \{N_d H(d, \mathbf{X}_i, Y_i) / N\pi_d\} \left\{ \sum_d \{N_d H(d, \mathbf{X}_i, Y_i) / (N\pi_d)\} \right\}^{-1}$ and remember that $\pi_0 + \pi_1 = 1$, we see that we can estimate π_0 by solving

$$\pi_0 = \sum_{i=1}^N H(0, \mathbf{X}_i, Y_i) \left[\sum_d N_d H(d, \mathbf{X}_i, Y_i) / \{\pi_0^{1-d} (1 - \pi_0)\} \pi_d \right]^{-1}.$$

ALGORITHM 1: Computing the Locally Efficient Score Function

The first two steps are done only once.

- Posit a model for $\eta_2(\epsilon, \mathbf{x})$ which has mean zero, and calculate (5), calling the result $S^*(\mathbf{X}, Y, D)$. Use $S^*(\cdot)$ in place of $S(\cdot)$ in (6)-(7).
- Estimate $f_{\mathbf{X}|D}(\mathbf{x}, d)$ by a kernel density estimate among the data with $D_i = d$, with result $\hat{f}_{\mathbf{X}|D}(\mathbf{x}, d)$.

The rest of the steps are done iteratively in the estimation algorithm.

- Solve $\hat{\pi}_0 = \sum_{i=1}^N H(0, \mathbf{X}_i, Y_i) \{N_0 H(0, \mathbf{X}_i, Y_i) / \hat{\pi}_0 + N_1 H(1, \mathbf{X}_i, Y_i) / (1 - \hat{\pi}_0)\}^{-1}$ to obtain $\hat{\pi}_0$ and set $\hat{\pi}_1 = 1 - \hat{\pi}_0$.
- In the definition of $\kappa(\mathbf{x}, y)$ in (6), form $\hat{\kappa}(\mathbf{x}, y)$ by replacing π_d by $\hat{\pi}_d$. Define $\hat{\kappa}_i = \hat{\kappa}(\mathbf{X}_i, Y_i)$.
- Define $\hat{f}_{di} = \hat{f}_{D|X,Y}(d, \mathbf{X}_i, Y_i) = N_d H(d, \mathbf{X}_i, Y_i) \hat{\kappa}_i / (N \hat{\pi}_d)$.
- For any function $h(d, \mathbf{x}, y)$ in (6), estimate $E\{h(D, \mathbf{X}, Y) \mid \mathbf{X}, D = d\}$ by nonparametric regression among observations with $D_i = d$.
- For any function $h(d, \mathbf{x}, y)$ in (6), estimate $E\{h(D, \mathbf{X}, Y) \mid D = d\}$ as $\hat{E}\{h(D, \mathbf{X}, Y) \mid D = d\} = \sum_{i=1}^N h(d, \mathbf{X}_i, Y_i) \hat{f}_{di} / \sum_{i=1}^N \hat{f}_{di}$.
- For any function $h(d, \mathbf{x}, y)$ in (6), estimate $E\{h(D, Y, \mathbf{X}) \mid \epsilon, \mathbf{X}\}$ by $\hat{E}\{h(D, Y, \mathbf{X}) \mid \epsilon, \mathbf{X}\} = \sum_{d=0}^1 N_d H(d, \mathbf{X}, Y) h(d, Y, \mathbf{X}) \hat{\kappa}(\mathbf{X}, Y) / (N \hat{\pi}_d)$.
- For any function $h(d, \mathbf{x}, y)$ in (6), estimate $E_{\text{true}}\{h(D, \mathbf{X}, Y) \mid \mathbf{X}\}$ by $\hat{E}_{\text{true}}\{h(D, \mathbf{X}, Y) \mid \mathbf{X}\} = \sum_{d=0}^1 \hat{\pi}_d \hat{E}\{h(d, \mathbf{X}, Y) \mid \mathbf{X}, D = d\} \hat{f}_{\mathbf{X}|D}(\mathbf{X}, d) / \sum_{d=0}^1 \hat{\pi}_d \hat{f}_{\mathbf{X}|D}(\mathbf{X}, d)$.

Application to the terms in (6) yields $\hat{g}(\epsilon_i, \mathbf{X}_i)$ and \hat{v}_d , and we then form $\hat{\mathbf{S}}_{\text{eff}}^*(D, \mathbf{X}_i, Y_i) = \mathbf{S}^*(\mathbf{X}, Y, D) - \hat{\mathbf{g}}(\epsilon, \mathbf{X}) - (1 - D)\hat{v}_0 - D\hat{v}_1$.

We have described the algorithm when \mathbf{X} is continuous. When \mathbf{X} is discrete, one simply replaces the density estimators and various nonparametric regressions with the corresponding averages associated with the different \mathbf{x} values.

- Next we have that

$$\begin{aligned} E(\mathbf{h} \mid \epsilon, \mathbf{X}) &= \sum_d \mathbf{h} f_{D|X,Y}(d, \mathbf{X}, Y) \\ &= \sum_d \{N_d H(d, \mathbf{X}, Y) \mathbf{h}(d, \mathbf{X}, Y) / (N \pi_d)\} \{ \sum_d N_d H(d, \mathbf{X}, Y) / (N \pi_d) \}^{-1}. \end{aligned}$$

- In addition,

$$\begin{aligned} E_{\text{true}}(\mathbf{h} \mid \mathbf{X}) &= \frac{\int \mathbf{h} \sum_d \pi_d f_{X,Y|D}^{\text{true}}(\mathbf{X}, y, d) d\mu(y)}{\int \sum_d \pi_d f_{X,Y|D}^{\text{true}}(\mathbf{X}, y, d) d\mu(y)} = \frac{\int \mathbf{h} \sum_d \pi_d f_{X,Y|D}(\mathbf{X}, y, d) d\mu(y)}{\int \sum_d \pi_d f_{X,Y|D}(\mathbf{X}, y, d) d\mu(y)} \\ &= \frac{\sum_d \pi_d \int \mathbf{h} f_{X,Y|D}(\mathbf{X}, y, d) d\mu(y)}{\sum_d \pi_d \int \mathbf{h} f_{Y|\mathbf{X},D}(\mathbf{X}, y, d) d\mu(y) f_{\mathbf{X}|D}(\mathbf{X}, d)} \\ &= \frac{\sum_d \pi_d \int \mathbf{h} f_{Y|\mathbf{X},D}(\mathbf{X}, y, d) d\mu(y) f_{\mathbf{X}|D}(\mathbf{X}, d)}{\sum_d \pi_d \int \mathbf{h} f_{Y|\mathbf{X},D}(\mathbf{X}, y, d) d\mu(y) f_{\mathbf{X}|D}(\mathbf{X}, d)} \\ &= \sum_d \pi_d E(\mathbf{h} \mid \mathbf{X}, d) f_{\mathbf{X}|D}(\mathbf{X}, d) / \sum_d \pi_d f_{\mathbf{X}|D}(\mathbf{X}, d), \end{aligned}$$

where in the last expression, both $f_{\mathbf{X}|D}(\mathbf{x}, d)$ and $E(\mathbf{h} \mid \mathbf{x}, d)$ need to be estimated nonparametrically.

- Finally, we have

$$E(\mathbf{h} \mid D = d) = \frac{\int f_{\mathbf{x},y}(\mathbf{x}, y) \mathbf{h}(d, \mathbf{x}, y) f_{D|X,Y}(d, \mathbf{x}, y) d\mu(\mathbf{x}) d\mu(y)}{\int f_{\mathbf{x},y}(\mathbf{x}, y) f_{D|X,Y}(d, \mathbf{x}, y) d\mu(\mathbf{x}) d\mu(y)},$$

which can be estimated as

$$\widehat{E}(\mathbf{h} \mid D = d) = \sum_{i=1}^N \mathbf{h}(d, \mathbf{X}_i, Y_i) f_{D|X,Y}(d, \mathbf{X}_i, Y_i) / \sum_{i=1}^N f_{D|X,Y}(d, \mathbf{X}_i, Y_i).$$

4.2 Distribution Theory

Because the locally efficient estimator is derived from well-established semiparametric procedures, while replacing the unknown quantities with nonparametric estimation in the proposed model, it is not surprising that it is asymptotically normally distributed with standard parametric rates of convergence. In addition, it achieves the semiparametric efficiency if the proposed model is correct. We describe the asymptotic properties of our estimator in Theorems 1, and provide a sketch of the proof for Theorem 1 in the Appendix. We first list the set of regularity conditions that Theorem 1 requires.

- C1: There exists constants $0 < C < \infty$ such that $\lim_{N \rightarrow \infty} N_1/N_2 = C$. In addition, the identifiability Assumptions 1 and 2 hold.
- C2: The univariate kernel function is a function that integrates to 1 and has support $(-1, 1)$ and order r , i.e., $\int K(x)x^t dx = 0$ if $1 \leq t < r$ and $\int K(x)x^r dx \neq 0$. The d -dimensional kernel function, still represented with K , is a product of d univariate kernel functions, that is, $K(\mathbf{x}) = \prod_{i=1}^d K(x_i)$ for a d -dimensional \mathbf{x} .
- C3: For $d = 1, 0$, $f_{X|D}(\mathbf{x} \mid D = d)$, $E(\epsilon^2 \kappa \mid \mathbf{X}, D = d)$, $E(\epsilon \mu_s \mid \mathbf{X}, D = d)$, $E(\epsilon f_0 \mid \mathbf{X}, D = d)$, $E(\epsilon f_1 \mid \mathbf{X}, D = d)$ have compact support and have continuous r^{th} derivatives.
- C4: The bandwidth $h = N^{-\tau}$ where $1/(2p) > \tau > 1/(4r)$, where p is the dimension of \mathbf{x} . This includes the optimal bandwidth $h = O(N^{-1/(2r+p)})$ as long as we choose a kernel of order $2r > p$.

Condition C1 ensures that there are a sufficient number of both cases and controls in the sample, which occurs in all case-control studies of the type we are studying (see the introductory paragraph). Conditions C2 and C4 are standard requirements on an r th order kernel function and on the bandwidth in the kernel smoothing literature (Ma and Zhu, 2013). Condition C3 is not the weakest possible. We impose this condition to simplify the technical proof. It can be replaced with weaker conditions in the region where $\|\mathbf{x}\|$ is large, at the expense of a more tedious technical treatment.

Theorem 1. We emphasize that for any random vector $\mathbf{S}(D, Y, X)$, expectation and covariance in the superpopulation is linked to expectation and covariance in the case-control sampling scheme (conditional on disease status) through

$$\begin{aligned} E\{\mathbf{S}(D, Y, X)\} &= \sum_{d=0}^1 (N_d/N) E\{\mathbf{S}(D, Y, X)|D = d\} \\ \text{cov}\{\mathbf{S}(D, Y, X)\} &= \sum_{d=0}^1 (N_d/N) \text{cov}\{\mathbf{S}(D, Y, X)|D = d\}. \end{aligned}$$

Under the regularity conditions C1-C4, in the case-control study, as $N \rightarrow \infty$, the estimator $\hat{\boldsymbol{\theta}}$ obtained from solving the estimating equation $\sum_{i=1}^N \hat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \hat{\boldsymbol{\theta}}) = 0$ satisfies

$$N^{1/2}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \rightarrow \text{Normal}\{\mathbf{0}, \mathbf{A}^{-1}\mathbf{B}(\mathbf{A}^{-1})^T\}$$

where $\mathbf{A} = E\{\partial \mathbf{S}_{\text{eff}}^*(D, \mathbf{X}, Y, \boldsymbol{\theta}_0)/\partial \boldsymbol{\theta}^T\}$ and $\mathbf{B} = \text{cov}\{\mathbf{S}_{\text{eff}}^*(D, \mathbf{X}, Y, \boldsymbol{\theta}_0)\}$.

5 Simulations

5.1 Setup

We performed a series of simulation studies in order to evaluate the finite sample performance of the various methods. In total, we considered 72 different cases. First, we considered a balanced design, where $N_0 = N_1 = 500$, and an imbalanced design with $N_0 = 666$ and $N_1 = 334$, i.e., 2 controls for every case. Second, we considered 3 disease rates: a relatively rare disease rate of 4.5%, an extremely rare disease rate of 0.5% and a common disease rate of 10%. The balanced design in rare or extremely rare disease cases is representative of a typical case-control study.

Third, we considered three settings for the logistic regression. We generated X from a Uniform(0,1) distribution. The logistic regression model was $\text{pr}(D = 1|Y, X) = H(\alpha_c + \alpha_1 X + \alpha_2 Y)$, where $\alpha_1 = 1$ and we varied $\alpha_2 = 0.00, 0.25, 0.50$. The regression model for Y given X is $Y = \beta_1 + \beta_2 X + \epsilon$, with $\beta_1 = 0$ and $\beta_2 = 1$.

Finally, we varied the distribution of the regression errors and whether they were homoscedastic or not, as follows.

- In the first set of simulations, we generated homoscedastic errors ϵ . The distribution of ϵ was either Normal(0, σ^2) with $\sigma^2 = 1$ or is a centered and standardized Gamma distribution with shape parameter 0.4, normalized to have mean zero and variance $\sigma^2 = 1$. To achieve an approximate 4.5% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-3.6, -3.8, -4.0)$. To achieve an approximate 0.5% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-5.8, -6.0, -6.2)$. To achieve an approximate 10% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-2.7, -2.9, -3.1)$.
- In the second set of simulations, we generated heteroscedastic errors as follows. The same distributions for ϵ were used, except that ϵ was multiplied by $(1 + X^2)^{3/4}/2$ in all the cases, so that $\text{var}(\epsilon|X) = (1 + X^2)^{3/2}/4$. To achieve an approximate 4.5% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-3.60, -3.75, -3.95)$. To achieve an approximate 0.5% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-5.8, -5.95, -6.2)$. To achieve an approximate 10% disease rate, for $\alpha_2 = (0.00, 0.25, 0.50)$ we set $\alpha_c = (-2.7, -2.9, -3.1)$.

With respect to the method described in Section 4.1, we mention the following details. The posited model η_2^* being a standard normal model in step 1. This yields the second component in S^* as $(y - \beta_1 - \beta_2 x)(1, x)^T$. In performing the many nonparametric calculations in steps 4, 5, 6, 7, we used a kernel estimates with a same bandwidth h throughout. We set the bandwidth at $h = cn_0^{-1/3}$, and experimented with different values c between $c = 0.5$ and $c = 2.0$, with little change in the results. To assess variability, we used the asymptotic results in Theorem 1, with the \mathbf{A} and \mathbf{B} matrices replaced by their corresponding sample averages evaluated at the estimated parameter values.

We compared our method with three others. The first was ordinary least squares among the controls, with sandwich standard errors: the sandwich method is used to adjust confidence intervals for possible heteroscedasticity. The second was the semiparametric efficient method that assumes normality and homoscedasticity, with standard errors obtained by inverting the Hessian of the loglikelihood (Lin and Zeng (2009)). The third was the method of Wei et al. (2013) that assumes homoscedasticity, but otherwise does not specify any particular error distribution model: we used the bootstrap to obtain standard errors for this method.

A striking conclusion of these simulations is that our methods, which assumes none of rare disease, normal errors or homoscedasticity, uniformly has coverage probabilities that achieve the nominal rates.

5.2 Homoscedastic Case

Results for the homoscedastic case are given in Tables 1-3. We display the mean estimate, the standard deviation across the simulations, the mean estimated standard deviation, coverage probabilities for nominal 90% and 95% confidence intervals, and the mean squared error efficiency of the methods relative to using only the controls.

The case $\alpha_2 = 0.00$ is interesting, because here Y is independent of D given X . Hence, all methods should achieve nominal coverage probabilities for estimating β , which is indeed seen in Table 1. Surprisingly, our method, which assumes neither normality nor homoscedasticity, is as efficient in terms of mean squared error as the semiparametric efficient method that assumes both, and is of course much more efficient than using only the controls.

For $\alpha_2 \neq 0$, and when ϵ is normally distributed, our method remains comparably as efficient as the semiparametric efficient method which assumes both normality and homoscedasticity. However, when the errors were not normally distributed, our method has much smaller bias and is much more efficient. In addition, the semiparametric efficient method has poor coverage probabilities when $\alpha_2 = 0.50$. While the method of Wei et al. (2013) maintains good coverage probabilities in all cases, our methods also maintains coverage, has smaller bias and is much more efficient.

5.3 Heteroscedastic Case

The results for the heteroscedastic case, with various disease rates and equal or unequal case-control ratios are given in Tables 4-6.

The results are much in line with the homoscedastic case, with a few important exceptions. The semiparametric efficient method, which assumes both homoscedasticity and normality, has a noticeable loss of coverage probability when $\alpha_2 \neq 0$, largely caused by bias. Because they used a bootstrap to compute standard errors, the method of Wei et al. (2013) maintains good coverage probability except when $\alpha_2 = 0.50$, where the bias causes deterioration in the coverage rates. Our method maintains good coverage probabilities in all cases, and because of its lack of bias, noticeably increased mean squared error efficiency.

6 Empirical Example

Epidemiological studies have led to the general belief that heterocyclic amines (HCA), such as MeIQx and PhIP, are significant risk factors associated with various forms of cancers, including colorectal cancer and breast cancer (Barrett et al., 2003; Sinha et al., 2001; De Stefani et al., 1997). One of the important food sources contributing to carcinogenic HCA, among many other potential sources, is red meat, which produces the agents during the cooking process. In addition, red meat contains other nutrients such as saturated fat which is also believed to relate to the occurrence of cancer. Due to this link, epidemiological and nutritional studies of cancer often include both red meat consumption and HCA as covariates to assess the risk of developing cancer, while simultaneously studying the relation between HCA amount and red meat consumption. Understanding this relation helps to understand the health impact of red meat consumption and is important in formulating food consumption guidelines for the general public.

We implemented our method on a data set involving colorectal adenoma, with 640 cases and 665 controls. The cases and controls were defined by the occurrence of colorectal adenoma (D). In our analysis, X is red meat consumption in grams. We used two different versions of Y , namely the heterocyclic amines MeIQx and PhIP that are produced during

the cooking of meat.

PhIP, MeIQx and red meat were transformed by adding 1.0 and taking logarithms to alleviate the heavy skewness of these measurements on the original scale. We also analyzed the subset of the study who were smokers. For the controls-only analysis, standard errors of the slope estimate were computed using the usual formula for least squares and also by the sandwich method. For our semiparametric analysis, we computed standard errors by the asymptotic formula of Theorem 1 and by the bootstrap, with 1,000 bootstrap samples. Given the results of the simulation, we do not expect any significant difference between these two estimates of standard errors for our method, with the asymptotic formula being much faster computationally.

We performed a preliminary analysis using only the controls. In the original data scale, all the covariates (PhIP, MeIQx and red meat consumption) are very skewed and heavy-tailed, see Figures S.1-S.2 in the **Supplementary Material**. The transformed data were much better behaved, see Figures S.3-S.4 in the **Supplementary Material**. Numerically, the skewness of MeIQx in the original and transformed data scales are 3.46 and -0.19, respectively. The skewness of PhIP in the original and transformed data scales are 7.93 and -0.20, respectively. Finally, the skewness of Red Meat in the original and transformed data scales are 1.78 and -0.58, respectively. These numbers and the plots indicate that the transformation did an acceptable to very good job of removing skewness.

Further preliminary analysis of the controls included scatterplots of the transformed data, both of which were reasonably well-behaved and indicated an increasing trend for increasing red meat consumption, consistent with a linear trend, see Figure S.5 in the **Supplementary Material**. To check this, we fit a quadratic model to the transformed data: in both cases, the p-value for the quadratic term exceeded 0.20, see Figure 2. Thus, we adopted a linear function for the mean $m(\cdot)$ in the subsequent secondary analysis. In addition, the regression of PhIP on red meat consumption is heavily heteroscedastic, while the regression of MeIQx on red meat is passably homoscedastic. This is shown in Figure 3, where we fit a regression of the absolute residuals from a quadratic fit against red meat consumption (Davidian and Carroll, 1987): the plots from a linear regression are essentially the same.

The results of this secondary analysis are given in Table 7. For MeIQx, the ordinary least

squares standard errors when using only the controls are roughly the same and that of the sandwich method, which makes sense since the regression is homoscedastic. In this case, as expected from the theory, our semiparametric approach has smaller standard errors, with the least squares standard errors being approximately 30% larger. For PhIP, where the regression is distinctly homoscedastic, the sandwich standard errors for ordinary least squares among the controls is roughly 30% larger than the standard error that assumes homoscedasticity, and roughly 40% larger than our semiparametric approach. As expected from the theory, where homoscedasticity is not assumed, the standard errors for our semiparametric approach are nearly the same using either the asymptotic formula or the bootstrap.

As a comparison, we also implemented the parametric method of Lin and Zeng (2009) as well as the robust method by Wei et al. (2013). Standard errors of the former were assessed both by using the inverse of the Hessian of the loglikelihood and by the bootstrap, while standard errors of the latter were assessed by the bootstrap alone. The parametric method's asymptotic standard error clearly under-estimates the variability for PhIP when compared to the bootstrap, something expected because of the heteroscedasticity in PhIP. For MeIQx, where the error is homoscedastic, the parametric method, the robust method and our semiparametric approach are almost identical.

In summary, in analyzing this data set, we verified the previous observation based on the control only data that the regression error from MeIQx and red meat consumption has homoscedastic error, while that from the PhIP and red meat consumption has heteroscedastic error. Our analysis also verified the positive relationship between red meat consumption and these two forms of HCA, indicating that increased red meat consumption leads to increased levels of MeIQx and PhIP, both being risk factors for colorectal cancer. The first order accuracy of the variability of the estimated slope for our method is validated though its near-identical result with the bootstrap, and of course through the simulation results.

7 Discussion

We have developed a locally efficient semiparametric estimator for the secondary analysis of case-control studies, where only a mean model is specified to describe the relationship

between the covariates. Despite this relatively weak assumption, we have shown that the problem is still identifiable under certain conditions. Through introducing the notion of a superpopulation, we are able to establish an estimation methodology via a conceptually tractable semiparametric procedure, although the derivation is highly non-standard and not trivial. The locally efficient estimator provides consistent estimation, and can achieve optimal efficiency if a posited regression error model happens to be true. Although the analysis is performed under the superpopulation concept, the general statements of consistency and local efficiency are valid in the case-control sampling scheme (Ma, 2010). In addition, the general methodology is applicable even if the linear logistic model (1) is replaced by other parametric models such as probit model, etc., as long as identifiability can be established.

Implementing the locally efficient estimator via Algorithm 1 requires several nonparametric regressions conditional on the covariates, which may be difficult when the dimension of the covariates increases. In such situations, dimension reduction techniques can be a good choice to achieve a balance between model flexibility and feasibility of parameter estimation and inference (Ma and Zhu, 2012). Further exploration of this is needed.

Acknowledgments

Ma's research was supported by NSF grant DMS-1206693 and NINDS grant R01-NS073671. Carroll's research was supported by National Cancer Institute grant R37-CA057030 and by NINDS grant R01-NS073671.

Appendix: Sketch of Technical Arguments

A.1 Proof of Proposition 1

Assume the contrary. That is, assume the problem is not identifiable. This means we can find parameters $\alpha_c, \boldsymbol{\alpha}_1, \alpha_2, \boldsymbol{\beta}, \eta_2, \eta_1$ and $\tilde{\alpha}_c, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2, \tilde{\boldsymbol{\beta}}, \tilde{\eta}_2, \tilde{\eta}_1$ so that, denoting $\tilde{\epsilon} = Y - m(\mathbf{x}, \tilde{\boldsymbol{\beta}})$,

$$\begin{aligned}\pi_d &= \int \eta_1(\mathbf{x})\eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} \frac{\exp\{d\alpha_c + du(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2,)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} d\mu(\mathbf{x})d\mu(y); \\ \tilde{\pi}_d &= \int \tilde{\eta}_1(\mathbf{x})\tilde{\eta}_2\{y - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}), \mathbf{x}\} \frac{\exp\{d\tilde{\alpha}_c + du(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)\}}{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \alpha_2)\}} d\mu(\mathbf{x})d\mu(y),\end{aligned}$$

we have that

$$\begin{aligned} & \frac{1}{\pi_d} \eta_1(\mathbf{x}) \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} \frac{\exp\{d\alpha_c + du(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \\ &= \frac{1}{\tilde{\pi}_d} \tilde{\eta}_1(\mathbf{x}) \tilde{\eta}_2\{y - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}), \mathbf{x}\} \frac{\exp\{d\tilde{\alpha}_c + du(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)\}}{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)\}} \end{aligned} \quad (\text{A.1})$$

for all (\mathbf{x}, y, d) . Take the ratio of the above expression at $d = 1$ and $d = 0$ respectively, we obtain that for all (\mathbf{x}, y) ,

$$\frac{\pi_0}{\pi_1} \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\} = \frac{\tilde{\pi}_0}{\tilde{\pi}_1} \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)\}.$$

This yields that $u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2) - u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)$ is a constant. Since it is zero at $(\mathbf{x}, y) = 0$, hence we have $u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2) - u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2) \equiv 0$. Thus, $\boldsymbol{\alpha}_1, \alpha_2 = \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2$, $\exp(\alpha_c)\pi_0/\pi_1 = \exp(\tilde{\alpha}_c)\tilde{\pi}_0/\tilde{\pi}_1$ and

$$\frac{1}{\pi_0} \frac{\eta_1(\mathbf{x}) \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} = \frac{1}{\tilde{\pi}_0} \frac{\tilde{\eta}_1(\mathbf{x}) \tilde{\eta}_2\{y - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}), \mathbf{x}\}}{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \tilde{\boldsymbol{\alpha}}_1, \tilde{\alpha}_2)\}}$$

for all (\mathbf{x}, y) . This gives

$$\tilde{\eta}_1(\mathbf{x}) \tilde{\eta}_2\{y - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}), \mathbf{x}\} = \frac{\tilde{\pi}_0}{\pi_0} \frac{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \eta_1(\mathbf{x}) \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\}. \quad (\text{A.2})$$

Integrating (A.2) and the product of (A.2) and y with respect to y , we obtain

$$\begin{aligned} \tilde{\eta}_1(\mathbf{x}) &= \frac{\tilde{\pi}_0}{\pi_0} \eta_1(\mathbf{x}) \int \frac{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} dy; \\ \tilde{\eta}_1(\mathbf{x}) m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) &= \frac{\tilde{\pi}_0}{\pi_0} \eta_1(\mathbf{x}) \int \frac{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} y dy \end{aligned}$$

respectively. Further taking ratios, we find

$$\begin{aligned} & \int \frac{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} y dy \\ &= m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) \int \frac{1 + \exp\{\tilde{\alpha}_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}}{1 + \exp\{\alpha_c + u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \alpha_2)\}} \eta_2\{y - m(\mathbf{x}, \boldsymbol{\beta}), \mathbf{x}\} dy. \end{aligned}$$

If $\alpha_c = \tilde{\alpha}_c$, then we obtain $m(\mathbf{x}, \boldsymbol{\beta}) = m(\mathbf{x}, \tilde{\boldsymbol{\beta}})$, hence $\boldsymbol{\beta} = \tilde{\boldsymbol{\beta}}$. We also obtain $\tilde{\eta}_1(\mathbf{x}) = \eta_1(\mathbf{x})\tilde{\pi}_0/\pi_0$. Since both $\tilde{\eta}_1(\mathbf{x})$ and $\eta_1(\mathbf{x})$ are valid density functions, we have $\tilde{\eta}_1(\mathbf{x}) = \eta_1(\mathbf{x})$ and $\pi_0 = \tilde{\pi}_0, \pi_1 = \tilde{\pi}_1$. This subsequently yields $\eta_2 = \tilde{\eta}_2$ contradicting our assumptions. Thus we obtain that $\alpha_c \neq \tilde{\alpha}_c$.

Denote

$$\begin{aligned} r(\epsilon, \mathbf{x}) &= \frac{1 + \exp[\tilde{\alpha}_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} \{\epsilon - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) + m(\mathbf{x}, \boldsymbol{\beta})\} \\ &= \exp(\tilde{\alpha}_c - \alpha_c) \{\epsilon - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) + m(\mathbf{x}, \boldsymbol{\beta})\} \\ &\quad + (1 - \exp(\tilde{\alpha}_c - \alpha_c)) \frac{\epsilon - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) + m(\mathbf{x}, \boldsymbol{\beta})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} \end{aligned}$$

By definition, η_2 is a valid conditional density function and it satisfies $\int \epsilon \eta_2(\epsilon, \mathbf{x}) d\epsilon = 0$, and we have that

$$\begin{aligned} 0 &= \int r(\epsilon, \mathbf{x}) \eta_2(\epsilon, \mathbf{x}) d\epsilon \\ &= -\exp(\tilde{\alpha}_c - \alpha_c) \{m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) - m(\mathbf{x}, \boldsymbol{\beta})\} \\ &\quad + (1 - \exp(\tilde{\alpha}_c - \alpha_c)) \int \frac{\epsilon - m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) + m(\mathbf{x}, \boldsymbol{\beta})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} \eta_2(\epsilon, \mathbf{x}) d\epsilon \end{aligned}$$

for all \mathbf{x} . This means

$$\begin{aligned} \frac{\{m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) - m(\mathbf{x}, \boldsymbol{\beta})\} \exp(\tilde{\alpha}_c - \alpha_c)}{1 - \exp(\tilde{\alpha}_c - \alpha_c)} &= \int \frac{\epsilon \eta_2(\epsilon, \mathbf{x})}{(1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}])} d\epsilon \\ &\quad - \int \frac{\{m(\mathbf{x}, \tilde{\boldsymbol{\beta}}) - m(\mathbf{x}, \boldsymbol{\beta})\} \eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \end{aligned}$$

for all \mathbf{x} . If we let $\mathbf{x} \rightarrow \mathbf{c}_1$, then

$$\begin{aligned} \frac{c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \exp(\tilde{\alpha}_c - \alpha_c)}{1 - \exp(\tilde{\alpha}_c - \alpha_c)} &= c_2 \int_{\mathcal{D}^c} \epsilon \eta_2(\epsilon, \mathbf{c}_1) d\epsilon - c_2 c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \int_{\mathcal{D}^c} \eta_2(\epsilon, \mathbf{c}_1) d\epsilon \\ &\quad + \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{\epsilon \eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \\ &\quad - \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \\ &= -c_2 c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) - c_2 \int_{\mathcal{D}} \epsilon \eta_2(\epsilon, \mathbf{c}_1) d\epsilon + c_2 c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \int_{\mathcal{D}} \eta_2(\epsilon, \mathbf{c}_1) d\epsilon \\ &\quad + \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{\epsilon \eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \\ &\quad - \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) \eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon. \end{aligned}$$

Thus,

$$\begin{aligned}
& \left| \frac{\exp(\tilde{\alpha}_c - \alpha_c)}{1 - \exp(\tilde{\alpha}_c - \alpha_c)} + c_2 \right| \\
= & \left| -c_2 \int_{\mathcal{D}} \frac{\epsilon \eta_2(\epsilon, \mathbf{c}_1)}{c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})} d\epsilon + c_2 \int_{\mathcal{D}} \eta_2(\epsilon, \mathbf{c}_1) d\epsilon \right. \\
& + \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{\epsilon \eta_2(\epsilon, \mathbf{x}) / c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \\
& \left. - \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} \int_{\mathcal{D}} \frac{\eta_2(\epsilon, \mathbf{x})}{1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}]} d\epsilon \right| \\
\leq & \frac{2}{|c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})|} \int_{\mathcal{D}} |\epsilon| \eta_2(\epsilon, \mathbf{c}_1) d\epsilon + 2 \int_{\mathcal{D}} \eta_2(\epsilon, \mathbf{c}_1) d\epsilon \\
\leq & \frac{2}{|c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})|} [E\{\epsilon^2 I(\epsilon \in \mathcal{D}) \mid \mathbf{c}_1\} \text{pr}(\epsilon \in \mathcal{D} \mid \mathbf{c}_1)]^{1/2} + 2 \text{pr}(\epsilon \in \mathcal{D} \mid \mathbf{c}_1) \\
\leq & \frac{2}{|c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})|} \{E(\epsilon^2) \delta\}^{1/2} + 2\delta.
\end{aligned}$$

We can make the upper bound of the above expression arbitrarily small by choosing δ arbitrarily close to zero, while the quantity on the left hand side is a constant. Hence we in fact have obtained

$$\frac{\exp(\tilde{\alpha}_c - \alpha_c)}{1 - \exp(\tilde{\alpha}_c - \alpha_c)} = -c_2$$

However, $-c_2$ is between -1 and 0 , simple calculation shows that these two constants cannot be equal, hence our problem is indeed identifiable.

A.2 Derivation of Λ and Λ^\perp

Consider the nuisance tangent space associated with η_1 and η_2 respectively, we have

$$\begin{aligned}
\Lambda_1 &= \{\mathbf{g}(\mathbf{x}) - E(\mathbf{g} \mid d) : \forall \mathbf{g} \text{ such that } E_{\text{true}}(\mathbf{g}) = \mathbf{0}\}; \\
\Lambda_2 &= \{\mathbf{g}(\epsilon, \mathbf{x}) - E(\mathbf{g} \mid d) : \forall \mathbf{g} \text{ such that } E_{\text{true}}(\mathbf{g} \mid \mathbf{X}) = E_{\text{true}}(\epsilon \mathbf{g} \mid \mathbf{X}) = \mathbf{0} \text{ a.s.}\}.
\end{aligned}$$

Hence $\Lambda = \Lambda_1 + \Lambda_2 = \{\mathbf{g}(\epsilon, \mathbf{x}) - E(\mathbf{g} \mid d) : \forall \mathbf{g} \text{ such that } E_{\text{true}}(\mathbf{g}) = E_{\text{true}}(\epsilon \mathbf{g} \mid \mathbf{X}) = \mathbf{0} \text{ a.s.}\}$. It is easily seen that $\Lambda_1^\perp = \{\mathbf{h} : E(\mathbf{h}) = \mathbf{0}, E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{X}\} = \mathbf{0} \text{ a.s.}\}$. This is because from

$$\begin{aligned}
0 &= E[\mathbf{h}^\top \{\mathbf{g}(\mathbf{X}) - E(\mathbf{g} \mid D)\}] \\
&= E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \{\mathbf{g}(\mathbf{X}) - E(\mathbf{g} \mid D)\}] \\
&= E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \mathbf{g}] \\
&= E(E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \mid \mathbf{X}] \mathbf{g}),
\end{aligned}$$

we obtain $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{X}\} \sum_d \int f_{X,Y,D}(\mathbf{X}, y, d) d\mu(y) / \eta_1(\mathbf{X}) = \mathbf{c}$ a.s. for some constant \mathbf{c} . Since $E[E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{X}\}] = \mathbf{0}$ a.s., we obtain

$$\mathbf{0} = \int E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{x}\} \sum_d \int f_{X,Y,D}(\mathbf{x}, y, d) d\mu(y) d\mu(\mathbf{x}) = \int \mathbf{c} \eta_1(\mathbf{x}) d\mu(\mathbf{x}) = \mathbf{c} \text{ a.s..}$$

Hence $\mathbf{c} = \mathbf{0}$ and $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{X}\} \sum_{d=0}^1 \int f_{X,Y,D}(\mathbf{X}, y, d) d\mu(y) / \eta_1(\mathbf{X}) = \mathbf{0}$ a.s., which yields $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \mathbf{X}\} = \mathbf{0}$ a.s..

Now we are in position to show

$$\begin{aligned} \Lambda^\perp &= \Lambda_1^\perp \cap \Lambda_2^\perp = [\mathbf{h}(d, \epsilon, \mathbf{x}) : E(\mathbf{h}) = \mathbf{0}, E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \\ &\quad \times \sum_d \frac{N_d}{N} \frac{H(d, \mathbf{X}, Y)}{p_D^{\text{true}}(d)} = \epsilon \mathbf{a}(\mathbf{X}) \text{ a.s.}], \end{aligned}$$

where $\mathbf{a}(\mathbf{x})$ is an arbitrary function of \mathbf{x} . This is because for any $\mathbf{h} \in \Lambda_1^\perp$, $\mathbf{h} \in \Lambda_2^\perp$ is equivalent to

$$\begin{aligned} 0 &= E[\mathbf{h}^\top \{\mathbf{g}(\epsilon, \mathbf{X}) - E(\mathbf{g} \mid D)\}] \\ &= E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \{\mathbf{g}(\epsilon, \mathbf{X}) - E(\mathbf{g} \mid D)\}] \\ &= E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \mathbf{g}] \\ &= E(E[\{\mathbf{h} - E(\mathbf{h} \mid D)\}^\top \mid \epsilon, \mathbf{X}] \mathbf{g}). \end{aligned}$$

Hence $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \sum_d f_{X,Y,D}(\mathbf{X}, Y, d) / \{\eta_1(\mathbf{X}) \eta_2(\epsilon, \mathbf{X})\} = \epsilon \mathbf{a}(\mathbf{X}) + \mathbf{c}(\mathbf{X})$ a.s.. Because $\mathbf{h} \in \Lambda_1^\perp$, we have $E[E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \mid \mathbf{X}] = \mathbf{0}$ a.s.. Hence

$$\begin{aligned} \mathbf{0} &= \int E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \frac{\sum_d f_{X,Y,D}(\mathbf{X}, y, d)}{\int \sum_d f_{X,Y,D}(\mathbf{X}, y, d) d\mu(y)} d\mu(y) \\ &= \frac{\int \{\epsilon \mathbf{a}(\mathbf{X}) + \mathbf{c}(\mathbf{X})\} \eta_1(\mathbf{x}) \eta_2(\epsilon, \mathbf{X}) d\mu(y)}{\int \sum_d f_{X,Y,D}(\mathbf{X}, y, d) d\mu(y)} \\ &= \frac{\mathbf{c}(\mathbf{X}) \eta_1(\mathbf{X})}{\int \sum_d f_{X,Y,D}(\mathbf{X}, y, d) d\mu(y)} \text{ a.s.,} \end{aligned}$$

hence $\mathbf{c}(\mathbf{X}) = \mathbf{0}$ a.s. and $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \sum_d f_{X,Y,D}(\mathbf{X}, Y, d) / \{\eta_1(\mathbf{X}) \eta_2(\epsilon, \mathbf{X})\} = \epsilon \mathbf{a}(\mathbf{X})$ a.s.. This means that $E\{\mathbf{h} - E(\mathbf{h} \mid D) \mid \epsilon, \mathbf{X}\} \sum_d (N_d/N) H(d, \mathbf{X}, Y) / p_D^{\text{true}}(d) = \epsilon \mathbf{a}(\mathbf{X})$ a.s..

A.3 Sketch of Proof of Theorem 1

For simplicity of proof, we split the N observations randomly into two sets. The first set contains $n_1 = N - N^{1-\delta}$ observations and the second set contains $n_2 = N^{1-\delta}$ observations, where δ is a small positive number. We form and solve the estimating equation using data in the first set, while calculating all the hatted quantities described in the algorithm using

data in the second set. We use this only as a technical device, although in our simulations and empirical example we used all the data.

In the algorithm, the approximations involve either replacing expectation with averaging, or standard kernel regression estimation or kernel density estimation, hence the differences between the quantities with hat and without hat have either mean zero, standard deviation $O(n_2^{-1/2})$, or mean $O(h^r)$, standard deviation $O\{(n_2h^p)^{-1/2}\}$. In particular, $\widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) - \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)$ has bias $O(h^r)$ and standard deviation $O\{(n_2h^p)^{-1/2}\}$. Recall the definition of expectation and covariance in the superpopulation explicitly written out in the statement of Theorem 1. Then

$$\begin{aligned}
\mathbf{0} &= n_1^{-1/2} \sum_{i=1}^{n_1} \widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \widehat{\boldsymbol{\theta}}) \\
&= n_1^{-1/2} \sum_{i=1}^{n_1} \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) + n_1^{-1/2} \sum_{i=1}^{n_1} \left\{ \widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) - \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) \right\} \\
&\quad + E \left\{ \frac{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}^T} + o_p(1) \right\} n_1^{1/2} (\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \\
&= n_1^{-1/2} \sum_{i=1}^{n_1} \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) + E \left\{ \frac{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}^T} \right\} n_1^{1/2} (\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \\
&\quad + n_1^{-1/2} \sum_{i=1}^{n_1} \left\{ \widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) - \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) \right\} + o_p(1).
\end{aligned}$$

We see that $\widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)$ differs from $\mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)$ in that all the unknown quantities, except \mathbf{S}^* , are estimated. This is equivalent to estimating the unknown functions $\eta_1(\mathbf{x})$, $\eta_2(\epsilon, \mathbf{x})$ in (4) and using the estimate $\widehat{\eta}_1(\mathbf{x})$, $\widehat{\eta}_2(\epsilon, \mathbf{x})$ in calculating $\mathbf{S}_{\text{eff}}^*$ from the posited \mathbf{S}^* . Thus, denoting $\widehat{\boldsymbol{\eta}} = (\widehat{\eta}_1, \widehat{\eta}_2)$, we can approximate

$$\begin{aligned}
&n_1^{-1/2} \sum_{i=1}^{n_1} \left\{ \widehat{\mathbf{S}}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) - \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) \right\} \\
&= n_1^{-1/2} \sum_{i=1}^{n_1} \left\{ \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \widehat{\boldsymbol{\eta}}) - \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \eta_0) \right\} \\
&= \left\{ n_1^{-1/2} \sum_{i=1}^{n_1} \partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \eta_0) / \partial \boldsymbol{\eta} \right\} (\widehat{\boldsymbol{\eta}} - \eta_0) + O_p\{n_1^{1/2} (\widehat{\boldsymbol{\eta}} - \eta_0)^2\} + o_p(1), \text{ (A.3)}
\end{aligned}$$

where $\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \eta_0) / \partial \boldsymbol{\eta}$ is pathwise derivative. However, $\mathbf{S}_{\text{eff}}^*$ is the projection of \mathbf{S}^*

to Λ^\perp so $\mathbf{S}_{\text{eff}}^* \in \Lambda^\perp$. Thus, for any parametric submodel of η involving parameter γ , we have

$$\begin{aligned} & E\{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \gamma) / \partial \gamma^T\} \\ &= \int \frac{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \gamma)}{\partial \gamma^T} f_{X,Y,D}(x, y, d) d\mu(x) \mu(y) d\mu(d) \\ &= - \int \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \gamma) \frac{\partial \log\{f_{X,Y,D}(x, y, d)\}}{\partial \gamma^T} f_{X,Y,D}(x, y, d) d\mu(x) \mu(y) d\mu(d) \\ &= -E\{\mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \gamma) \mathbf{S}_\gamma^T\} = \mathbf{0}. \end{aligned}$$

The last equality is because by definition $\mathbf{S}_\gamma \in \Lambda$ which is orthogonal to Λ^\perp and $\mathbf{S}_{\text{eff}}^* \in \Lambda^\perp$. Here, $f_{X,Y,D}(x, y, d)$ is defined in (4). Because γ is parameter of any arbitrary submodel of η , we actually have obtained

$$E\{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \eta_0) / \partial \eta\} = -E\{\mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0, \eta_0) \mathbf{S}_\eta^T\} = \mathbf{0},$$

where \mathbf{S}_η is the nuisance score function along the arbitrarily chosen specific path of the pathwise derivative. Thus, the first term of (A.3) is of order $o_p(1)$. On the other hand, $O_p\{n_1^{1/2}(\widehat{\eta} - \eta_0)^2\} = O_p\{n_1^{1/2}h^{2r} + n_1^{1/2}(n_2h^p)^{-1}\} = o_p(1)$. We therefore obtain

$$\mathbf{0} = n_1^{-1/2} \sum_{i=1}^{n_1} \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0) + E\left\{\frac{\partial \mathbf{S}_{\text{eff}}^*(D_i, \mathbf{X}_i, Y_i, \boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}^T}\right\} n_1^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + o_p(1).$$

This yields $n_1^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \rightarrow \text{Normal}\{\mathbf{0}, \mathbf{A}^{-1}\mathbf{B}(\mathbf{A}^{-1})^T\}$, and hence

$$N^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \rightarrow \text{Normal}\{\mathbf{0}, \mathbf{A}^{-1}\mathbf{B}(\mathbf{A}^{-1})^T\}$$

when $N \rightarrow \infty$.

References

- Barrett, J. H., Smith, G., Waxman, R., Gooderham, N., Lightfoot, T., Garner, R. C., Augustsson, K., Wolf, C. R., Bishop, D. T., Forman, D., and the Colorectal Cancer Study Group (2003). Investigation of interaction between n-acetyltransferase 2 and heterocyclic amines as potential risk factors for colorectal cancer. *Carcinogenesis*, 24, 275–282.
- Bickel, P. J., Klassen, C. A. J., Ritov, Y., and Wellner, J. A. (1993). *Efficient and Adaptive Estimation for Semiparametric Models*. Johns Hopkins University Press, Baltimore.
- Chatterjee, N. and Carroll, R. J. (2005). Semiparametric maximum likelihood estimation in case-control studies of gene-environment interactions. *Biometrika*, 92, 399–418.
- Chatterjee, N., Chen, J., Spinka, C., and Carroll, R. J. (2006). Comment on the paper Likelihood based inference on haplotype effects in genetic association studies by D. Y. Lin and D. Zeng. *Journal of the American Statistical Association*, 101, 108–110.

- Chen, Y. H., Chatterjee, N., and Carroll, R. J. (2008). Retrospective analysis of haplotype-based case-control studies under a flexible model for gene-environment association. *Biostatistics*, 9, 81–99.
- Chen, Y. H., Chatterjee, N., and Carroll, R. J. (2009). Shrinkage estimators for robust and efficient inference in haplotype-based case-control studies. *Journal of the American Statistical Association*, 104, 220–233.
- Corley, D. A., Jensen, C. D., Marks, A. R., Zhao, W. K., Lee, J. K., Doubeni, C. A., Zauber, A. G., de Boer, J., Fireman, B. H., Schottinger, J. E., Schottinger, M., Q., V. P., G., N. R., L., T. R., and Q., C. P. (2014). Adenoma detection rate and risk of colorectal cancer and death. *New England Journal of Medicine*, 370, 1298–1306.
- Davidian, M. and Carroll, R. J. (1987). Variance function estimation. *Journal of the American Statistical Association*, 82, 1079–1092.
- De Stefani, E., Ronco, A., Mendilaharsu, M., Guidobono, M., and Deneo-Pellegrini, H. (1997). Meat intake, heterocyclic amines, and risk of breast cancer: a case-control study in uruguay. *Cancer Epidemiology Biomarkers & Prevention*, 6, 573–581.
- He, J., Li, H., Edmondson, A. C., Rader, D. J., and Li, M. (2012). A gaussian copula approach for the analysis of secondary phenotypes in case-control genetic association studies. *Biostatistics*, 13, 497–508.
- Jiang, Y., Scott, A. J., and Wild, C. J. (2006). Secondary analysis of case-control data. *Statistics in Medicine*, 25, 1323–1339.
- Li, H., Gail, M. H., Berndt, S., and Chatterjee, N. (2010). Using cases to strengthen inference on the association between single nucleotide polymorphisms and a secondary phenotype in genome-wide association studies. *Genetic Epidemiology*, 34, 427–433.
- Lin, D. Y. and Zeng, D. (2009). Proper analysis of secondary phenotype data in case-control association studies. *Genetic Epidemiology*, 33, 256–265.
- Lobach, I., Carroll, R. J., Spinka, C., Gail, M. H., and Chatterjee, N. (2008). Haplotype-based regression analysis of case-control studies with unphased genotypes and measurement errors in environmental exposures. *Biometrics*, 64, 673–684.
- Ma, Y. (2010). A semiparametric efficient estimator in case-control studies. *Bernoulli*, 16, 585–603.
- Ma, Y. and Zhu, L. P. (2012). A semiparametric approach to dimension reduction. *Journal of the American Statistical Association*, 107, 168–179.
- Ma, Y. and Zhu, L. P. (2013). Efficient estimation in sufficient dimension reduction. *Annals of Statistics*, 41, 250–268.
- Prentice, R. L. and Pyke, R. (1979). Logistic disease incidence models and case-control studies. *Biometrika*, 66, 403–411.

- Scott, A. J. and Wild, C. J. (2002). On the robustness of weighted methods for fitting models to case-control data. *Journal of the Royal Statistical Society, Series B*, 64, 207–219.
- Sinha, R., Kulldorff, M., Chow, W.-H., Denobile, J., and Rothman, N. (2001). Dietary intake of heterocyclic amines, meat-derived mutagenic activity, and risk of colorectal adenomas. *Cancer Epidemiology Biomarkers & Prevention*, 10, 559–562.
- Tsiatis, A. A. (2006). *Semiparametric Theory and Missing Data*. Springer, New York.
- Wei, J., Carroll, R. J., Muller, U., Van Keilegom, I., and Chatterjee, N. (2013). Locally efficient estimation for homoscedastic regression in the secondary analysis of case-control data. *Journal of the Royal Statistical Society, Series B*, 75, 185–206.
- Yamaji, Y., Mitsushima, T., Ikuma, H., Watabe, H., Okamoto, M., Kawabe, T., Wada, R., Doi, H., and Omata, M. (2004). Incidence and recurrence rates of colorectal adenomas estimated by annually repeated colonoscopies on asymptomatic japanese. *Gut*, 53, 568–572.




Figure 1: Illustration of the bias induced by the case-control sampling scheme. The red solid line is the true regression function, while the blue dashed line is the regression function when using all the data and ignoring the case-control sampling scheme.




Figure 2: The fitted curves from a quadratic regression of MeIQx (solid red line) and PhIP (dashed blue line) on red meat consumption, using the controls. The fitted values were normalized to fit on the same plot. Neither have a statistically significant quadratic term.




Figure 3: Plots to diagnose heteroscedasticity, with the curves representing relative standard deviation as a function of red meat consumption. Plotted are the fitted curves from a linear regression of the absolute residuals of the regression of MeIQx (solid red line) and PhIP (dashed blue line) on red meat consumption, using the controls. The fitted values were normalized to be equal at the minimum value of red meat consumption. The essentially flat curve for MeIQx indicates homoscedasticity, while that for PhIP is very strongly heteroscedastic. The latter has implications for data analysis, see Table 7 and the discussion in Section 6.

	Normal				Gamma			
$\alpha_2 = 0.00$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.998	0.998	1.001	1.008	0.996	1.000	0.997	1.001
s.d.	0.151	0.110	0.114	0.109	0.155	0.110	0.120	0.110
Est. sd	0.155	0.110	0.122	0.130	0.154	0.110	0.122	0.116
90%	0.903	0.900	0.921	0.910	0.898	0.894	0.910	0.912
95%	0.952	0.955	0.957	0.959	0.958	0.954	0.956	0.959
MSE Eff		1.878	1.734	1.909		1.966	1.663	1.987
$\alpha_2 = 0.25$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.980	0.983	0.976	0.998	0.977	0.962	0.961	0.993
s.d.	0.151	0.113	0.116	0.113	0.151	0.139	0.115	0.093
Est. sd	0.154	0.111	0.119	0.115	0.148	0.140	0.120	0.103
90%	0.906	0.878	0.895	0.900	0.895	0.902	0.895	0.912
95%	0.947	0.939	0.953	0.966	0.939	0.948	0.943	0.963
MSE Eff		1.785	1.663	1.816		1.129	1.599	2.682
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.974	0.969	0.946	0.992	0.954	0.799	0.958	1.002
s.d.	0.146	0.106	0.119	0.116	0.139	0.179	0.133	0.099
Est. sd	0.154	0.112	0.122	0.126	0.139	0.173	0.132	0.103
90%	0.918	0.909	0.884	0.915	0.885	0.681	0.892	0.917
95%	0.961	0.955	0.943	0.964	0.934	0.787	0.943	0.961
MSE Eff		1.780	1.270	1.627		0.295	1.092	2.186

Table 1: Results of the simulation study with $n_1 = 500$ cases and $n_0 = 500$ controls, disease rate of approximately 4.5%, with homoscedastic errors. Here "Normal" means that $\epsilon = \text{Normal}(0, 1)$, while "Gamma" means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only ("Controls"), the semiparametric efficient method that assumes normality and homoscedasticity ("Param"), the method of Wei, et al. (2012), ("Robust"), and our method ("Semi"). Over 1,000 simulations, we computed the mean estimated β ("Mean"), its standard deviation ("s.d."), the mean estimated standard deviation ("Est. sd"), the coverage for a nominal 90% confidence interval ("90%"), the coverage for a nominal 95% confidence interval ("95%"), and the mean squared error efficiency compared to using only the controls ("MSE Eff").

		Normal				Gamma			
disease rate 10%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.913	0.876	0.784	0.979	0.885	0.885	0.929	0.993	
s.d.	0.120	0.121	0.159	0.117	0.124	0.124	0.108	0.109	
Est. sd	0.119	0.123	0.154	0.117	0.153	0.126	0.110	0.109	
90%	0.806	0.746	0.600	0.893	0.870	0.792	0.847	0.897	
95%	0.867	0.837	0.723	0.956	0.926	0.891	0.908	0.948	
MSE Eff		0.731	0.305	1.554		0.951	1.628	2.279	
disease rate 0.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.991	0.996	0.987	1.010	0.978	0.854	1.029	0.991	
s.d.	0.165	0.114	0.118	0.121	0.148	0.231	0.155	0.097	
Est. sd	0.155	0.112	0.120	0.122	0.149	0.223	0.160	0.096	
90%	0.876	0.893	0.904	0.898	0.902	0.830	0.904	0.895	
95%	0.925	0.942	0.949	0.938	0.945	0.904	0.950	0.945	
MSE Eff		2.099	1.938	1.852		0.300	0.900	2.359	

Table 2: Results of the simulation study with $n_1 = 500$ cases and $n_0 = 500$ controls, $\alpha_2 = 0.5$, homoscedastic errors. Here “Normal” means that $\epsilon = \text{Normal}(0, 1)$, while “Gamma” means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only (“Controls”), the semiparametric efficient method that assumes normality and homoscedasticity (“Param”), the method of Wei, et al. (2012), (“Robust”), and our method (“Semi”). Over 1,000 simulations, we computed the mean estimated β (“Mean”), its standard deviation (“s.d.”), the mean estimated standard deviation (“Est. sd”), the coverage for a nominal 90% confidence interval (“90%”), the coverage for a nominal 95% confidence interval (“95%”), and the mean squared error efficiency compared to using only the controls (“MSE Eff”).

		Normal				Gamma			
disease rate 4.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.962	0.960	0.956	0.994	0.951	0.856	0.936	0.996	
s.d.	0.133	0.106	0.108	0.113	0.128	0.153	0.123	0.101	
Est. sd	0.133	0.110	0.113	0.121	0.120	0.152	0.120	0.108	
90%	0.892	0.884	0.893	0.901	0.845	0.751	0.844	0.916	
95%	0.957	0.943	0.954	0.952	0.925	0.848	0.910	0.960	
MSE Eff		1.491	1.407	1.494		0.426	0.977	1.839	
disease rate 10%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.921	0.850	0.831	0.991	0.937	0.879	0.927	1.060	
s.d.	0.106	0.114	0.134	0.082	0.129	0.117	0.107	0.082	
Est. sd	0.103	0.113	0.136	0.080	0.133	0.117	0.110	0.077	
90%	0.797	0.621	0.673	0.900	0.872	0.739	0.840	0.908	
95%	0.881	0.752	0.780	0.949	0.932	0.845	0.909	0.949	
MSE Eff		0.492	0.375	2.568		0.727	1.228	1.996	
disease rate 0.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	1	0.997	0.991	1.004	0.997	0.901	1.018	1.000	
s.d.	0.133	0.107	0.113	0.110	0.129	0.191	0.134	0.100	
Est. sd	0.134	0.111	0.111	0.113	0.130	0.190	0.142	0.099	
90%	0.904	0.911	0.894	0.904	0.890	0.858	0.925	0.897	
95%	0.944	0.959	0.943	0.945	0.947	0.921	0.966	0.953	
MSE Eff		1.544	1.377	1.460		0.360	0.911	1.665	

Table 3: Results of the simulation study with $n_1 = 334$ cases and $n_0 = 666$ controls, $\alpha_2 = 0.5$, homoscedastic errors. Here “Normal” means that $\epsilon = \text{Normal}(0, 1)$, while “Gamma” means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only (“Controls”), the semiparametric efficient method that assumes normality and homoscedasticity (“Param”), the method of Wei, et al. (2012), (“Robust”), and our method (“Semi”). Over 1,000 simulations, we computed the mean estimated β (“Mean”), its standard deviation (“s.d.”), the mean estimated standard deviation (“Est. sd”), the coverage for a nominal 90% confidence interval (“90%”), the coverage for a nominal 95% confidence interval (“95%”), and the mean squared error efficiency compared to using only the controls (“MSE Eff”).

	Normal				Gamma			
$\alpha_2 = 0.00$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.996	0.996	1.000	1.005	0.992	0.994	1.000	1.002
s.d.	0.099	0.071	0.071	0.076	0.099	0.070	0.073	0.077
Est. sd	0.096	0.070	0.072	0.082	0.096	0.070	0.071	0.078
90%	0.887	0.892	0.895	0.898	0.887	0.903	0.893	0.898
95%	0.932	0.953	0.949	0.950	0.944	0.946	0.947	0.951
MSE Eff		1.948	1.961	1.692		1.971	1.847	1.663
$\alpha_2 = 0.25$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.986	1.044	0.973	0.997	0.983	1.063	0.964	0.995
s.d.	0.100	0.072	0.066	0.077	0.094	0.082	0.069	0.071
Est. sd	0.096	0.071	0.070	0.081	0.094	0.083	0.072	0.074
90%	0.880	0.838	0.907	0.912	0.894	0.825	0.863	0.904
95%	0.936	0.907	0.953	0.959	0.946	0.900	0.934	0.950
MSE Eff		1.415	1.984	1.717		0.852	1.516	1.801
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi
Mean	0.972	1.088	0.949	0.991	0.962	1.145	0.906	0.993
s.d.	0.099	0.072	0.068	0.083	0.095	0.096	0.076	0.082
Est. sd	0.096	0.072	0.071	0.102	0.090	0.100	0.076	0.105
90%	0.877	0.664	0.842	0.897	0.857	0.591	0.655	0.900
95%	0.936	0.789	0.914	0.946	0.909	0.714	0.756	0.935
MSE Eff		0.816	1.479	1.519		0.343	0.717	1.546

Table 4: Results of the simulation study with $n_1 = 500$ cases and $n_0 = 500$ controls, disease rate of approximately 4.5%, with heteroscedastic errors. Here "Normal" means that $\epsilon = \text{Normal}(0, 1)$, while "Gamma" means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only ("Controls"), the semiparametric efficient method that assumes normality and homoscedasticity ("Param"), the method of Wei, et al. (2012), ("Robust"), and our method ("Semi"). Over 1,000 simulations, we computed the mean estimated β ("Mean"), its standard deviation ("s.d."), the mean estimated standard deviation ("Est. sd"), the coverage for a nominal 90% confidence interval ("90%"), the coverage for a nominal 95% confidence interval ("95%"), and the mean squared error efficiency compared to using only the controls ("MSE Eff").

		Normal				Gamma			
disease rate 10%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.905	0.897	1.078	0.990	0.950	0.931	1.065	1.001	
s.d.	0.083	0.073	0.091	0.117	0.101	0.073	0.071	0.108	
Est. sd	0.083	0.072	0.089	0.115	0.100	0.072	0.072	0.111	
90%	0.676	0.600	0.770	0.895	0.847	0.765	0.781	0.895	
95%	0.766	0.698	0.850	0.947	0.914	0.851	0.859	0.955	
MSE Eff		0.998	1.107	1.154		1.258	1.370	1.089	
disease rate 0.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.997	1.113	0.973	1.007	0.991	1.296	0.890	0.995	
s.d.	0.098	0.073	0.067	0.082	0.102	0.113	0.087	0.072	
Est. sd	0.101	0.072	0.070	0.088	0.098	0.112	0.084	0.071	
90%	0.906	0.541	0.892	0.897	0.895	0.145	0.630	0.907	
95%	0.951	0.663	0.957	0.942	0.937	0.231	0.745	0.941	
MSE Eff		0.531	1.842	1.419		0.104	0.533	2.013	

Table 5: Results of the simulation study with $n_1 = 500$ cases and $n_0 = 500$ controls, $\alpha_2 = 0.5$, heteroscedastic errors. Here “Normal” means that $\epsilon = \text{Normal}(0, 1)$, while “Gamma” means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only (“Controls”), the semiparametric efficient method that assumes normality and homoscedasticity (“Param”), the method of Wei, et al. (2012), (“Robust”), and our method (“Semi”). Over 1,000 simulations, we computed the mean estimated β (“Mean”), its standard deviation (“s.d.”), the mean estimated standard deviation (“Est. sd”), the coverage for a nominal 90% confidence interval (“90%”), the coverage for a nominal 95% confidence interval (“95%”), and the mean squared error efficiency compared to using only the controls (“MSE Eff”).

		Normal				Gamma			
disease rate 4.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.977	1.052	0.961	0.996	0.961	1.085	0.926	0.994	
s.d.	0.084	0.070	0.063	0.077	0.083	0.087	0.066	0.082	
Est. sd	0.087	0.072	0.064	0.083	0.08	0.087	0.067	0.090	
90%	0.883	0.825	0.859	0.913	0.827	0.735	0.702	0.918	
95%	0.939	0.892	0.930	0.952	0.905	0.831	0.806	0.954	
MSE Eff		0.998	1.382	1.276		0.568	0.855	1.244	
disease rate 10%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.911	0.909	1.021	1.001	0.956	0.937	1.027	1.000	
s.d.	0.072	0.064	0.080	0.079	0.084	0.066	0.070	0.087	
Est. sd	0.072	0.065	0.080	0.076	0.087	0.065	0.072	0.094	
90%	0.654	0.595	0.895	0.901	0.867	0.772	0.877	0.906	
95%	0.749	0.700	0.949	0.951	0.927	0.851	0.933	0.952	
MSE Eff		1.058	1.915	2.099		1.080	1.597	1.188	
disease rate 0.5%									
$\alpha_2 = 0.50$	Controls	Param	Robust	Semi	Controls	Param	Robust	Semi	
Mean	0.997	1.073	0.979	1.007	0.994	1.189	0.920	0.997	
s.d.	0.088	0.073	0.066	0.078	0.084	0.100	0.071	0.070	
Est. sd	0.087	0.072	0.063	0.086	0.085	0.099	0.073	0.069	
90%	0.891	0.728	0.871	0.901	0.899	0.384	0.725	0.911	
95%	0.950	0.820	0.929	0.952	0.953	0.539	0.829	0.960	
MSE Eff		0.727	1.616	1.264		0.155	0.620	1.445	

Table 6: Results of the simulation study with $n_1 = 334$ cases and $n_0 = 666$ controls, $\alpha_2 = 0.5$, heteroscedastic errors. Here “Normal” means that $\epsilon = \text{Normal}(0, 1)$, while “Gamma” means that ϵ is a centered and scale Gamma random variable with shape 0.4, mean zero and variance one. The analyses performed were using controls only (“Controls”), the semiparametric efficient method that assumes normality and homoscedasticity (“Param”), the method of Wei, et al. (2012), (“Robust”), and our method (“Semi”). Over 1,000 simulations, we computed the mean estimated β (“Mean”), its standard deviation (“s.d.”), the mean estimated standard deviation (“Est. sd”), the coverage for a nominal 90% confidence interval (“90%”), the coverage for a nominal 95% confidence interval (“95%”), and the mean squared error efficiency compared to using only the controls (“MSE Eff”).

All Data						
	Controls only			Estimate	Parametric	
	Estimate	OLS se	Sandwich se		Asymptotic se	Bootstrap se
MeIQx	0.868	0.034	0.035	0.862	0.026	0.026
PhIP	0.742	0.064	0.080	0.751	0.046	0.056
	Robust			Estimate	Semiparametric	
	Estimate	Bootstrap se			Asymptotic se	Bootstrap se
MeIQx	0.862	0.028		0.862	0.027	0.027
PhIP	0.751	0.057		0.750	0.057	0.058
Smokers only						
	Controls only			Estimate	Parametric	
	Estimate	OLS se	Sandwich se		Asymptotic se	Bootstrap se
MeIQx	0.816	0.050	0.057	0.847	0.036	0.037
PhIP	0.619	0.095	0.132	0.737	0.063	0.080
	Robust			Estimate	Semiparametric	
	Estimate	Bootstrap se			Asymptotic se	Bootstrap se
MeIQx	0.847	0.038		0.846	0.036	0.039
PhIP	0.737	0.084		0.736	0.082	0.087

Table 7: Results of data analysis when Y is either MeIQx or PhIP. For the controls only, “OLS se” is the ordinary least squares standard error estimate, while “Sandwich se” is the sandwich method standard error estimate. For the parametric and semiparametric analysis, “Asymptotic se” is the standard error estimate from asymptotic theory, while “Bootstrap se” is the bootstrap standard error. For the robust analysis, only bootstrap standard error is available. The regression of PhIP on red meat (X) is heteroscedastic, reflected in the difference between the OLS standard error and the Sandwich standard error for the controls only analysis, as well as the difference between the asymptotic standard error and the bootstrap standard error of the parametric estimator.

Supplementary Material for *Semiparametric Estimation in Secondary Analysis of Case-Control Studies*

Yanyuan Ma and Raymond J. Carroll

Department of Statistics, 3143 TAMU, Texas A&M University, College Station, TX

77843-3143

ma@stat.tamu.edu and carroll@stat.tamu.edu

S.1 Derivation of the Efficient Score

Having obtained both the score function and the two spaces Λ and Λ^\perp , we only need to project the score function onto Λ^\perp to obtain the efficient score \mathbf{S}_{eff} . To do this, we write $\mathbf{S}_\theta = \mathbf{S} - E(\mathbf{S} | D) = \mathbf{g}(\epsilon, \mathbf{X}) - E(\mathbf{g} | D) + \mathbf{S}_{\text{eff}}$, where $E_{\text{true}}(\mathbf{g}) = E_{\text{true}}(\epsilon \mathbf{g} | \mathbf{X}) = \mathbf{0}$. We alternatively write $\mathbf{S}_{\text{eff}} = \mathbf{S} - \mathbf{g}(\epsilon, \mathbf{X}) - E(\mathbf{S} - \mathbf{g} | D)$ and \mathbf{S}_{eff} satisfies $E\{\mathbf{S}_{\text{eff}} - E(\mathbf{S}_{\text{eff}} | D) | \epsilon, \mathbf{X}\} \sum_d (N_d/N) H(d, \mathbf{x}, Y) / p_D^{\text{true}}(d) = \epsilon \mathbf{a}(\mathbf{X})$ and $E(\mathbf{S}_{\text{eff}}) = 0$. However, $E(\mathbf{S}_{\text{eff}} | d) = 0$ automatically, hence we can ignore the second requirement and the first requirement simplifies to $E(\mathbf{S}_{\text{eff}} | \epsilon, \mathbf{X}) \sum_d (N_d/N) \{H(d, \mathbf{X}, Y) / p_D^{\text{true}}(d)\} = \epsilon \mathbf{a}(\mathbf{X})$. This gives

$$\epsilon \mathbf{a}(\mathbf{X}) \left\{ \sum_d \frac{N_d H(d, \mathbf{X}, Y)}{N p_D^{\text{true}}(d)} \right\}^{-1} = E(\mathbf{S} - \mathbf{g} | \epsilon, \mathbf{X}) - E\{E(\mathbf{S} - \mathbf{g} | D) | \epsilon, \mathbf{X}\}.$$

It follows that

$$f_{D|X,Y}(d, \mathbf{x}, y) = \frac{N_d H(d, \mathbf{x}, y)}{N p_D^{\text{true}}(d)} \left\{ \sum_d \frac{N_d H(d, \mathbf{x}, y)}{N p_D^{\text{true}}(d)} \right\}^{-1}.$$

To simplify notation, in the following calculation we denote

$$\begin{aligned}
\pi_0 &= p_D^{\text{true}}(0) = \int \eta_1(\mathbf{x})\eta_2(\epsilon, \mathbf{x})H(0, \mathbf{x}, y)d\mu(\mathbf{x})d\mu(y); \\
\pi_1 &= p_D^{\text{true}}(1) = \int \eta_1(\mathbf{x})\eta_2(\epsilon, \mathbf{x})H(1, \mathbf{x}, y)d\mu(\mathbf{x})d\mu(y); \\
b_0 &= E\{f_{D|\mathbf{X},Y}(1, \mathbf{X}, Y) \mid D = 0\}; \\
b_1 &= E\{f_{D|\mathbf{X},Y}(0, \mathbf{X}, Y) \mid D = 1\}; \\
\mathbf{c}_0 &= E(\mathbf{S} \mid D = 0) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 0\}; \\
\mathbf{c}_1 &= E(\mathbf{S} \mid D = 1) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 1\}; \\
\kappa(\mathbf{x}, y) &= [\sum_{d=0}^1 \{N_d H(d, \mathbf{x}, y)\} / (N\pi_d)]^{-1}; \\
\mathbf{u}_0 &= E\{\epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0\}; \\
\mathbf{u}_1 &= E\{\epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 1\}; \\
\mathbf{v}_0 &= E(\mathbf{S} - \mathbf{g} \mid D = 0); \\
\mathbf{v}_1 &= E(\mathbf{S} - \mathbf{g} \mid D = 1).
\end{aligned}$$

Note that $\pi_0 + \pi_1 = 1$, $b_0 N_0 = b_1 N_1$, $\mathbf{c}_0 N_0 + \mathbf{c}_1 N_1 = \mathbf{0}$ and $\mathbf{v}_0 \pi_0 + \mathbf{v}_1 \pi_1 = \mathbf{0}$.

Under a true model, $\pi_0, \pi_1, b_0, b_1, \mathbf{c}_0, \mathbf{c}_1$ are known quantities, while $\mathbf{u}_1, \mathbf{u}_1, \mathbf{v}_0, \mathbf{v}_1$ are not known because $\mathbf{g} = \mathbf{g}(\epsilon, \mathbf{x})$ and $\mathbf{a} = \mathbf{a}(\mathbf{x})$ are not specified. To further obtain $\mathbf{u}_1, \mathbf{u}_1, \mathbf{v}_0, \mathbf{v}_1$,

$$\begin{aligned}
\epsilon \mathbf{a}(\mathbf{x})\kappa(\mathbf{x}, y) &= E(\mathbf{S} - \mathbf{g} \mid \epsilon, \mathbf{X} = \mathbf{x}) - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{x}, y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{x}, y) \\
&= E(\mathbf{S} \mid \epsilon, \mathbf{X} = \mathbf{x}) - \mathbf{g} - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{x}, y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{x}, y).
\end{aligned}$$

Alternatively, we also have

$$\mathbf{g}(\epsilon, \mathbf{x}) = E(\mathbf{S} \mid \epsilon, \mathbf{X} = \mathbf{x}) - \epsilon \mathbf{a}(\mathbf{x})\kappa(\mathbf{x}, y) - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{x}, y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{x}, y). \quad (\text{S.1})$$

Since $\mathbf{v}_0 = E(\mathbf{S} - \mathbf{g} \mid D = 0)$, we obtain

$$\begin{aligned}
\mathbf{v}_0 &= E(\mathbf{S} \mid D = 0) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) - \epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \\
&\quad - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{X}, Y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{X}, Y) \mid D = 0\} \\
&= \mathbf{c}_0 + \mathbf{u}_0 + \mathbf{v}_0(1 - b_0) + \mathbf{v}_1 b_0.
\end{aligned}$$

Thus, we have $b_0 \mathbf{v}_0 - b_0 \mathbf{v}_1 - \mathbf{u}_0 = \mathbf{c}_0$. Similarly, from $\mathbf{v}_1 = E(\mathbf{S} - \mathbf{g} \mid D = 1)$, we obtain

$$\begin{aligned}
\mathbf{v}_1 &= E(\mathbf{S} \mid D = 1) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) - \epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \\
&\quad - \mathbf{v}_0 f_{D|X,Y}(0, \mathbf{X}, Y) - \mathbf{v}_1 f_{D|X,Y}(1, \mathbf{X}, Y) \mid D = 1\} \\
&= \mathbf{c}_1 + \mathbf{u}_1 + \mathbf{v}_0 b_1 + \mathbf{v}_1(1 - b_1).
\end{aligned}$$

Thus, we have $-b_1\mathbf{v}_0 + b_1\mathbf{v}_1 - \mathbf{u}_1 = \mathbf{c}_1$. Since

$$E\{\epsilon\mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y)\} = \mathbf{0},$$

we have $\mathbf{u}_0N_0 + \mathbf{u}_1N_1 = \mathbf{0}$. Since $E_{\text{true}}(\mathbf{S} - \mathbf{g}) = \mathbf{0}$, we have

$$\begin{aligned} \mathbf{0} &= \sum_d \int (\mathbf{S} - \mathbf{g}) f_{\mathbf{X}, Y|D}(\mathbf{x}, y, d) \{f_{X, Y, D}^{\text{true}}(\mathbf{x}, y, d) / f_{\mathbf{X}, Y|D}(\mathbf{x}, y, d)\} d\mu(\mathbf{x}) d\mu(y) \\ &= \sum_d \int (\mathbf{S} - \mathbf{g}) f_{\mathbf{X}, Y|D}(\mathbf{x}, y, d) \{f_{X, Y, D}^{\text{true}}(\mathbf{x}, y, d) / f_{\mathbf{X}, Y|D}^{\text{true}}(\mathbf{x}, y, d)\} d\mu(\mathbf{x}) d\mu(y) \\ &= \sum_d \int (\mathbf{S} - \mathbf{g}) f_{\mathbf{X}, Y|D}(\mathbf{x}, y, d) p_D^{\text{true}}(d) d\mu(\mathbf{x}) d\mu(y) \\ &= \pi_0\mathbf{v}_0 + \pi_1\mathbf{v}_1. \end{aligned}$$

Combining the above relations, we have obtained $N_0\mathbf{u}_0 + N_1\mathbf{u}_1 = \mathbf{0}$, $\pi_0\mathbf{v}_0 + \pi_1\mathbf{v}_1 = \mathbf{0}$, $b_0\mathbf{v}_0 - b_0\mathbf{v}_1 - \mathbf{u}_0 = \mathbf{c}_0$ and $-b_1\mathbf{v}_0 + b_1\mathbf{v}_1 - \mathbf{u}_1 = \mathbf{c}_1$. The last two equations are equivalent so one is redundant. Using these relations, we can rewrite $\mathbf{u}_1, \mathbf{v}_0, \mathbf{v}_1$ as a function of \mathbf{u}_0 :

$$\mathbf{u}_1 = -(N_0/N_1)\mathbf{u}_0, \quad \mathbf{v}_0 = (\pi_1/b_0)(\mathbf{u}_0 + \mathbf{c}_0), \quad \mathbf{v}_1 = -(\pi_0/b_0)(\mathbf{u}_0 + \mathbf{c}_0). \quad (\text{S.2})$$

We cannot obtain a more explicit expression for \mathbf{u}_0 at this stage, but we can further obtain $\mathbf{a}(\mathbf{x})$ as a function of \mathbf{u}_0 . Using (S.1) and since $E_{\text{true}}(\epsilon\mathbf{g} | \mathbf{X}) = \mathbf{0}$, we have

$$\begin{aligned} E_{\text{true}}\{\epsilon E(\mathbf{S} | \epsilon, \mathbf{X}) | \mathbf{X}\} - E_{\text{true}}\{\epsilon^2\kappa(\mathbf{X}, Y) | \mathbf{X}\} \mathbf{a}(\mathbf{X}) \\ - \mathbf{v}_0 E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\} - \mathbf{v}_1 E_{\text{true}}\{\epsilon f_{D|X, Y}(1, \mathbf{X}, Y) | \mathbf{X}\} = \mathbf{0}. \end{aligned}$$

Hence

$$\begin{aligned} \mathbf{a}(\mathbf{X}) &= [E_{\text{true}}\{\epsilon^2\kappa(\mathbf{X}, Y) | \mathbf{X}\}]^{-1} \\ &\quad [E_{\text{true}}\{\epsilon E(\mathbf{S} | \epsilon, \mathbf{X}) | \mathbf{X}\} - \mathbf{v}_0 E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\} \\ &\quad \quad - \mathbf{v}_1 E_{\text{true}}\{\epsilon f_{D|X, Y}(1, \mathbf{X}, Y) | \mathbf{X}\}] \\ &= [E_{\text{true}}\{\epsilon^2\kappa(\mathbf{X}, Y) | \mathbf{X}\}]^{-1} [E_{\text{true}}\{\epsilon E(\mathbf{S} | \epsilon, \mathbf{X}) | \mathbf{X}\} \\ &\quad - (\pi_1/b_0)(\mathbf{u}_0 + \mathbf{c}_0) E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\} \\ &\quad + (\pi_0/b_0)(\mathbf{u}_0 + \mathbf{c}_0) E_{\text{true}}\{\epsilon f_{D|X, Y}(1, \mathbf{X}, Y) | \mathbf{X}\}]. \end{aligned}$$

To further simplify notation, denote

$$\begin{aligned} t_1(\mathbf{X}) &= [E_{\text{true}}\{\epsilon^2\kappa(\mathbf{X}, Y) | \mathbf{X}\}]^{-1}; \\ \mathbf{t}_2(\mathbf{X}) &= E_{\text{true}}\{\epsilon E(\mathbf{S} | \epsilon, \mathbf{X}) | \mathbf{X}\} - (\pi_1/b_0)\mathbf{c}_0 E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\} \\ &\quad + (\pi_0/b_0)\mathbf{c}_0 E_{\text{true}}\{\epsilon f_{D|X, Y}(1, \mathbf{X}, Y) | \mathbf{X}\}; \\ &= E_{\text{true}}\{\epsilon E(\mathbf{S} | \epsilon, \mathbf{X}) | \mathbf{X}\} - (\mathbf{c}_0/b_0) E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\}; \\ t_3(\mathbf{X}) &= -(\pi_1/b_0) E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\} + (\pi_0/b_0) E_{\text{true}}\{\epsilon f_{D|X, Y}(1, \mathbf{X}, Y) | \mathbf{X}\} \\ &= -b_0^{-1} E_{\text{true}}\{\epsilon f_{D|X, Y}(0, \mathbf{X}, Y) | \mathbf{X}\}. \end{aligned} \quad (\text{S.3})$$

Then

$$\mathbf{a}(\mathbf{x}) = t_1(\mathbf{x})\{\mathbf{t}_2(\mathbf{x}) + t_3(\mathbf{x})\mathbf{u}_0\}, \quad (\text{S.4})$$

hence the definition of \mathbf{u}_0 yields

$$\begin{aligned} \mathbf{u}_0 &= E[\epsilon\{t_1(\mathbf{X})\{\mathbf{t}_2(\mathbf{X}) + t_3(\mathbf{X})\mathbf{u}_0\}\}\kappa(\mathbf{X}, Y) \mid D = 0] \\ &= E[\epsilon t_1(\mathbf{X})\mathbf{t}_2(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0] + E[\epsilon t_1(\mathbf{X})t_3(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0]\mathbf{u}_0. \end{aligned}$$

This yields

$$\mathbf{u}_0 = (1 - E[\epsilon t_1(\mathbf{X})t_3(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0])^{-1} E[\epsilon t_1(\mathbf{X})\mathbf{t}_2(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0]. \quad (\text{S.5})$$

Combining the above results, we have obtained the analytic form of $\mathbf{S}_{\text{eff}} = \mathbf{S} - \mathbf{g} - E(\mathbf{S} - \mathbf{g} \mid D = d)$, where \mathbf{g} is given in (S.1), $\mathbf{a}(\mathbf{x})$ is given in (S.4), $\mathbf{v}_0, \mathbf{v}_1$ are given in (S.2) \mathbf{u}_0 is given in (S.5) and the functions t_1, \mathbf{t}_2, t_3 are given in (S.3).

In forming the estimating equation $\sum_{i=1}^N \mathbf{S}_{\text{eff}} = 0$, we will have $\sum_{i=1}^N [\mathbf{S}(\mathbf{X}_i, Y_i, D_i) - \mathbf{g}\{Y_i - m(\mathbf{X}_i, \boldsymbol{\beta}), \mathbf{X}_i\}] - N_0 E(\mathbf{S} - \mathbf{g} \mid D = 0) - N_1 E(\mathbf{S} - \mathbf{g} \mid D = 1) = \mathbf{0}$. Using (S.1), we obtain

$$\begin{aligned} E(\mathbf{S} - \mathbf{g} \mid D = 0) &= E(\mathbf{S} \mid D = 0) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 0\} + E\{\epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 0\} \\ &\quad + \mathbf{v}_0 E\{f_{D|X,Y}(0, \mathbf{X}, Y) \mid D = 0\} + \mathbf{v}_1 E\{f_{D|X,Y}(1, \mathbf{X}, Y) \mid D = 0\} \\ &= \mathbf{c}_0 + \mathbf{u}_0 + \mathbf{v}_0(1 - b_0) + \mathbf{v}_1 b_0 \end{aligned}$$

and

$$\begin{aligned} E(\mathbf{S} - \mathbf{g} \mid D = 1) &= E(\mathbf{S} \mid D = 1) - E\{E(\mathbf{S} \mid \epsilon, \mathbf{X}) \mid D = 1\} + E\{\epsilon \mathbf{a}(\mathbf{X})\kappa(\mathbf{X}, Y) \mid D = 1\} \\ &\quad + \mathbf{v}_0 E\{f_{D|X,Y}(0, \mathbf{X}, Y) \mid D = 1\} + \mathbf{v}_1 E\{f_{D|X,Y}(1, \mathbf{X}, Y) \mid D = 1\} \\ &= \mathbf{c}_1 + \mathbf{u}_1 + \mathbf{v}_0 b_1 + \mathbf{v}_1(1 - b_1), \end{aligned}$$

hence

$$\begin{aligned} &N_0 E(\mathbf{S} - \mathbf{g} \mid D = 0) + N_1 E(\mathbf{S} - \mathbf{g} \mid D = 1) \\ &N_0 \{\mathbf{c}_0 + \mathbf{u}_0 + \mathbf{v}_0(1 - b_0) + \mathbf{v}_1 b_0\} + N_1 \{\mathbf{c}_1 + \mathbf{u}_1 + \mathbf{v}_0 b_1 + \mathbf{v}_1(1 - b_1)\} \\ &(N_0 \mathbf{c}_0 + N_1 \mathbf{c}_1) + (N_0 \mathbf{u}_0 + N_1 \mathbf{u}_1) + (N_0 \mathbf{v}_0 + N_1 \mathbf{v}_1) + (\mathbf{v}_1 - \mathbf{v}_0)(N_0 b_0 - N_1 b_1). \end{aligned}$$

Thus, the estimating equation simplifies to $\sum_{i=1}^N [\mathbf{S}(\mathbf{X}_i, Y_i, D_i) - \mathbf{g}\{Y_i - m(\mathbf{X}_i, \boldsymbol{\beta}), \mathbf{X}_i\}] - N_0 \mathbf{v}_0 - N_1 \mathbf{v}_1 = \mathbf{0}$.

S.2 Verification of Assumptions 1-2 in the Linear Model as in Remark 2

Let $\mathcal{D}^c = [-K, K]$ for a sufficiently large K . We have assumed that the conditional distribution of ϵ satisfies the property that $\text{pr}(|\epsilon| > K | \mathbf{X} = \mathbf{x}) \rightarrow 0$ as $K \rightarrow \infty$ uniformly in \mathbf{x} , and hence $\text{pr}(\epsilon \in \mathcal{D})$ can be made arbitrarily small uniformly in \mathbf{x} .

Without loss of generality, assume the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is not zero. We consider three situations, 1. $\tilde{\beta}_c - \beta_c \neq 0$, 2. $\tilde{\beta}_c - \beta_c = 0$ and the first component of $\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1$ is not zero and 3) $\tilde{\beta}_c - \beta_c = 0$ and the first component of $\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1$ is zero. Note that ϵ has conditional mean zero. Thus for most common density functions with bounded variance function, this requirement is satisfied.

Case 1 and Case 2: If the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is positive, let $\mathbf{c}_1 = (\infty, \mathbf{0}^T)^T$, otherwise, let $\mathbf{c}_1 = (-\infty, \mathbf{0}^T)^T$. Let $c_2 = 0$, $\mathcal{D}^c = [-K, K]$ for a sufficiently large K . Then

$$\begin{aligned} & \sup_{\epsilon \in \mathcal{D}^c} \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} |(1 + \exp[\alpha_c + u\{\mathbf{x}, m(\mathbf{x}, \boldsymbol{\beta}) + \epsilon, \boldsymbol{\alpha}_1, \alpha_2\}])^{-1} - c_2| \\ &= \sup_{-K \leq \epsilon \leq K} \lim_{\mathbf{x} \rightarrow \mathbf{c}_1} |[1 + \exp\{\alpha_c + \mathbf{x}^T(\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2) + \beta_c \alpha_2 + \epsilon \alpha_2\}]^{-1}| \\ &= 0. \end{aligned}$$

Finally, since $\mathcal{D} = (-\infty, -K) \cup (K, \infty)$, we have that every element $u \in \mathcal{D}$ satisfies $u > 1$ as long as $K > 1$. We have thus verified Assumption 1.

We have $c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) = \mathbf{c}_1^T(\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1) + (\tilde{\beta}_c - \beta_c)$, which is not zero if $\tilde{\beta}_c - \beta_c \neq 0$ (case 1), and is also not zero if the first component of $\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1$ is not zero (case 2). Thus, Assumption 2 also holds.

Case 3: Since $\tilde{\boldsymbol{\beta}} \neq \boldsymbol{\beta}$, without loss of generality, we assume the second component of $\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1$ is not zero in this case. We select \mathbf{c}_1 as follows. If the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is positive and the second component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is non-negative, let $\mathbf{c}_1 = (\infty, \infty, \mathbf{0}^T)^T$. If the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is positive and the second component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is negative, let $\mathbf{c}_1 = (\infty, -\infty, \mathbf{0}^T)^T$. If the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is negative and the second component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is non-negative, let $\mathbf{c}_1 = (-\infty, \infty, \mathbf{0}^T)^T$. If the first component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is negative and the second component of $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2$ is negative, let $\mathbf{c}_1 = (-\infty, -\infty, \mathbf{0}^T)^T$. The selection of c_2, \mathcal{D}, K remains the same as in Cases 1 and 2. We can see the same arguments lead to the verification of Assumption 1. In addition, $c(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}}) = \mathbf{c}_1^T(\tilde{\boldsymbol{\beta}}_1 - \boldsymbol{\beta}_1)$, which is either ∞ or $-\infty$, and is thus not zero. Thus, Assumption 2 also holds.

S.3 Verification of Nonidentifiability in the Special Case in Remark 2

Here, we provide the details of the proof of the nonidentifiability result in Remark 2, which happens when when $m(\mathbf{x}, \boldsymbol{\beta}) = \mathbf{x}^T \boldsymbol{\beta}_1 + \beta_c$ and $u(\mathbf{x}, y, \boldsymbol{\alpha}_1, \boldsymbol{\alpha}_c) = \mathbf{x}^T \boldsymbol{\alpha}_1 + y \alpha_2 + \alpha_c$, and $\boldsymbol{\alpha}_1 + \boldsymbol{\beta}_1 \alpha_2 = \mathbf{0}$. We first exclude a special case when $\alpha_2 = 0$. This special case implies $\boldsymbol{\alpha}_1 = \mathbf{0}$ and $\alpha_2 = 0$, hence the case-control sampling is in fact random sampling. Thus, in the following, we assume $\alpha_2 \neq 0$. We point out that

$$m_1 \equiv \int \epsilon \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) \neq 0.$$

This is because we can use the mean value theorem for integration to obtain

$$\begin{aligned} & \int \epsilon \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) \\ &= \int_{-\infty}^0 \epsilon \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) + \int_0^{\infty} \epsilon \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) \\ &= \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 - k_1 \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 - k_1 \alpha_2)} \int_{-\infty}^0 \epsilon \eta_2(\epsilon) d\mu(\epsilon) + \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + k_2 \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + k_2 \alpha_2)} \int_0^{\infty} \epsilon \eta_2(\epsilon) d\mu(\epsilon) \\ &= \left\{ \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + k_2 \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + k_2 \alpha_2)} - \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 - k_1 \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 - k_1 \alpha_2)} \right\} \int_0^{\infty} \epsilon \eta_2(\epsilon) d\mu(\epsilon) \\ &= \frac{\{\exp(\tilde{\alpha}_c) - \exp(\alpha_c)\} \exp(\beta_c \alpha_2) \{\exp(k_2 \alpha_2) - \exp(-k_1 \alpha_2)\}}{\{1 + \exp(\alpha_c + \beta_c \alpha_2 + k_2 \alpha_2)\} \{1 + \exp(\alpha_c + \beta_c \alpha_2 - k_1 \alpha_2)\}} \int_0^{\infty} \epsilon \eta_2(\epsilon) d\mu(\epsilon) \neq 0, \end{aligned}$$

where k_1, k_2 are positive constants.

Following the notation in the proof of Proposition 1, we define $\eta_2(\epsilon, \mathbf{x}) = \eta_2(\epsilon)$,

$$\begin{aligned} \tilde{\eta}_1(\mathbf{x}) &= \frac{\tilde{\pi}_0}{\pi_0} \eta_1(x) \int \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon); \\ \tilde{\beta}_c &= \beta_c + \left\{ \int \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) \right\}^{-1} m_1; \\ \tilde{\eta}_2(\epsilon) &= c_0 \frac{1 + \exp(\tilde{\alpha}_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon + \tilde{\beta}_c - \beta_c), \end{aligned}$$

where

$$c_0^{-1} = \int \frac{1 + \exp(\tilde{\alpha}_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon + \tilde{\beta}_c - \beta_c) d\mu(\epsilon).$$

Obviously $\int \tilde{\eta}_2(\epsilon) d\mu(\epsilon) = 1$. We can easily verify that

$$\begin{aligned} \int \tilde{\eta}_1(\mathbf{x}) d\mu(\mathbf{x}) &= \frac{\tilde{\pi}_0}{\pi_0} \int \eta_1(x) \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon) d\mu(\epsilon) d\mu(\mathbf{x}) \\ &= \frac{\tilde{\pi}_0}{\pi_0} \{ \pi_0 + \exp(\tilde{\alpha}_c - \alpha_c) \pi_1 \} \\ &= \tilde{\pi}_0 + \frac{\tilde{\pi}_0 \exp(\tilde{\alpha}_c) \pi_1}{\pi_0 \exp(\alpha_c)} = \tilde{\pi}_0 + \tilde{\pi}_1 = 1, \end{aligned}$$

using the intermediate results in the proof of Proposition 1. We can also obtain

$$\begin{aligned} \int \epsilon \tilde{\eta}_2(\epsilon) d\mu(\epsilon) &= c_0 \int \epsilon \frac{1 + \exp(\tilde{\alpha}_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)}{1 + \exp(\alpha_c + \tilde{\beta}_c \alpha_2 + \epsilon \alpha_2)} \eta_2(\epsilon + \tilde{\beta}_c - \beta_c) d\mu(\epsilon) \\ &= c_0 \int (t + \beta_c - \tilde{\beta}_c) \frac{1 + \exp(\tilde{\alpha}_c + \beta_c \alpha_2 + t \alpha_2)}{1 + \exp(\alpha_c + \beta_c \alpha_2 + t \alpha_2)} \eta_2(t) d\mu(\epsilon) = 0. \end{aligned}$$

Now we can easily verify that the two parameter sets $\{\alpha_c, \boldsymbol{\alpha}_1, \alpha_2, \boldsymbol{\beta}_1, \beta_c, \eta_1(\mathbf{x}), \eta_2(\epsilon)\}$ and $\{\tilde{\alpha}_c, \boldsymbol{\alpha}_1, \alpha_2, \boldsymbol{\beta}_1, \tilde{\beta}_c, \tilde{\eta}_1(\mathbf{x}), \tilde{\eta}_2(\epsilon)\}$ satisfy (A.1), hence the problem is not identifiable.




Figure S.1: Boxplots of the variables in the original data scale among the controls. Each variable has been normalized to have maximum value 1.0. This and Figure S.2 indicate a need for data transformation.




Figure S.2: Kernel density estimates in the original data scale, among the controls. This and Figure S.1 indicate a need for data transformation.




Figure S.3: Boxplots of the variables in the transformed data scale, among the controls. Each variable has been normalized to have maximum value 1.0. Contrast with the boxplots in the original data scale in Figure S.1.




Figure S.4: Kernel density estimates in the transformed data scale, among the controls. Contrast with the boxplots in the original data scale in Figure S.2





Figure S.5: Scatterplots of transformed MeIQX and PhIP against transformed Red Meat, among the controls.