

Formula Sheets – post-Test 2 material – STAT 515

Test statistic (hypothesis test about μ):

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

test statistic for μ_D in paired-sample t -test: Same as above but with \bar{D} and s_D .

Test statistic (hypothesis test about p):

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

CI for $\mu_1 - \mu_2$:

$$(\bar{x}_1 - \bar{x}_2) \pm t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

Test statistic (comparing two means, independent samples):

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

CI for $p_1 - p_2$:

$$(\hat{p}_1 - \hat{p}_2) \pm z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$$

Test statistic (comparing two proportions):

$$z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

where \hat{p} is the pooled sample proportion.

ANOVA table formulas:

$$MST = SST/(p-1), \quad MSE = SSE/(n-p), \quad F = MST/MSE$$

Regression and correlation formulas:

$$\hat{\beta}_1 = SS_{xy}/SS_{xx}, \quad \hat{\beta}_0 = \bar{Y} - \hat{\beta}_1\bar{X}, \quad s = \sqrt{MSE} = \sqrt{SSE/(n-2)},$$

$$r = \frac{SS_{xy}}{\sqrt{SS_{xx}SS_{yy}}}, \quad r^2 = 1 - SSE/SS_{yy}$$

Test statistic for test of model usefulness:

$$t = \frac{\hat{\beta}_1}{s/\sqrt{SS_{xx}}}$$

CI for β_1 :

$$\hat{\beta}_1 \pm t_{\alpha/2}(s/\sqrt{SS_{xx}})$$

CI for $E(Y)$ at $x = x_p$:

$$\hat{Y} \pm t_{\alpha/2}(s)\sqrt{1/n + (x_p - \bar{x})^2/SS_{xx}}$$

PI for new Y at $x = x_p$:

$$\hat{Y} \pm t_{\alpha/2}(s)\sqrt{1 + 1/n + (x_p - \bar{x})^2/SS_{xx}}$$

Test statistic, Test for Multinomial Probabilities:

$$\chi^2 = \sum \frac{[n_i - E(n_i)]^2}{E(n_i)}$$

where $E(n_i) = np_i$ is the expected cell count if H_0 is true.

Test statistic, Test for Independence:

$$\chi^2 = \sum \frac{[n_{ij} - \hat{E}(n_{ij})]^2}{\hat{E}(n_{ij})}$$

where $\hat{E}(n_{ij}) = r_i c_j / n$ is the expected count in cell (i, j) under independence.