

# STAT 513 fa 2019 Exam II

Karl B. Gregory

*Do not open this test until told to do so; no calculators allowed; no notes allowed; no books allowed; show your work so that partial credit may be given.*

Some upper quantiles of some  $t$ -distributions:

$\xi$	0.10	0.05	0.025	0.01	0.005
$t_{28,\xi}$	1.3125	1.7011	2.0484	2.4671	2.7633
$t_{29,\xi}$	1.3114	1.6991	2.0452	2.4620	2.7564
$t_{30,\xi}$	1.3104	1.6973	2.0423	2.4573	2.7500
$t_{31,\xi}$	1.3095	1.6955	2.0395	2.4528	2.7440

Some upper quantiles of some chi-squared distributions:

$\xi$	0.10	0.05	0.025	0.01	0.005
$\chi_{1,\xi}^2$	2.71	3.84	5.02	6.63	7.88
$\chi_{2,\xi}^2$	4.61	5.99	7.38	9.21	10.60

For any  $\mathbf{a} \in \mathbb{R}^{p+1}$ , a  $(1 - \alpha)100\%$  confidence interval for  $\mathbf{a}^T \hat{\boldsymbol{\beta}}$  is given by

$$\mathbf{a}^T \hat{\boldsymbol{\beta}} \pm t_{n-p-1, \alpha/2} \hat{\sigma} \sqrt{\mathbf{a}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{a}},$$

and the  $t$ -test statistic for comparing  $\mathbf{a}^T \boldsymbol{\beta}$  to a null value  $a^*$  is given by

$$T_n = \frac{\mathbf{a}^T \hat{\boldsymbol{\beta}} - a^*}{\hat{\sigma} \sqrt{\mathbf{a}^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{a}}}.$$

Let  $X_1, \dots, X_n$  be a random sample with likelihood function  $L(\theta; X_1, \dots, X_n)$ . Then for hypotheses of the form  $H_0: \theta = \theta_0$  versus  $H_1: \theta \neq \theta_0$  the likelihood ratio is

$$\text{LR}(X_1, \dots, X_n) = \frac{L(\theta_0; X_1, \dots, X_n)}{L(\hat{\theta}; X_1, \dots, X_n)},$$

where  $\hat{\theta}$  is the MLE of  $\theta$ .

1. Each of  $n$  randomly selected subjects is given  $m$  opportunities to predict the outcome of a coin toss. Let  $p$  denote the probability that a subject makes the correct prediction. The numbers of correct predictions  $Y_1, \dots, Y_n$  made by the subjects are assumed to be independent  $\text{Binomial}(m, p)$  random variables. Researchers from L'Institut de la Clairvoyance wish to test the hypotheses

$$H_0: p = 1/2 \text{ versus } H_1: p \neq 1/2.$$

The likelihood function is given by

$$L(p; Y_1, \dots, Y_n) = \prod_{i=1}^n \binom{m}{Y_i} p^{Y_i} (1-p)^{m-Y_i}.$$

- Write down the log-likelihood function  $\ell(p; Y_1, \dots, Y_n)$ . It will be convenient to use  $\sum_{i=1}^n Y_i = n\bar{Y}_n$ .
  - Find the maximum likelihood estimator of  $p$ .
  - Write down the likelihood ratio  $\text{LR}(Y_1, \dots, Y_n)$  and simplify it as much as seems reasonable.
  - Give the form of the likelihood ratio test for the hypotheses of interest.
  - Get an expression for  $-2 \log \text{LR}(Y_1, \dots, Y_n)$  and state the rejection rule of the asymptotic likelihood ratio test of size  $\alpha = 0.05$ .
2. Consider the simple linear regression model

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

for the data  $(Y_1, x_1), \dots, (Y_n, x_n)$ , where  $\varepsilon_1, \dots, \varepsilon_n$  are independent  $\text{Normal}(0, \sigma^2)$  error terms and  $x_1, \dots, x_n$  are fixed constants. Suppose we will reject  $H_0: \beta_1 = 0$  in favor of  $H_1: \beta_1 \neq 0$  if and only if

$$\frac{|\hat{\beta}_1|}{\hat{\sigma} \sqrt{S_{xx}^{-1}}} > t_{n-2, \alpha/2},$$

where  $\hat{\beta}_1$  is the least-squares estimator of  $\beta_1$  and  $S_{xx} = \sum_{i=1}^n (x_i - \bar{x}_n)^2$ . Fill in the blanks.

- A \_\_\_\_\_ (larger/smaller) error term variance leads to \_\_\_\_\_ (larger/smaller) power of the test when  $\beta_1 \neq 0$ .
  - Values of  $x_1, \dots, x_n$  with a \_\_\_\_\_ (larger/smaller) spread lead to \_\_\_\_\_ (larger/smaller) power of the test when  $\beta_1 \neq 0$ .
  - Values of  $\beta_1$  of \_\_\_\_\_ (larger/smaller) magnitude lead to \_\_\_\_\_ (larger/smaller) power of the test.
  - A \_\_\_\_\_ (larger/smaller) value of  $\alpha$  leads to \_\_\_\_\_ (larger/smaller) power of the test when  $\beta_1 \neq 0$ .
3. Suppose  $\text{Var}(\hat{\beta}_0) = 2$ ,  $\text{Var}(\hat{\beta}_1) = 1$ , and  $\text{Cov}(\hat{\beta}_0, \hat{\beta}_1) = -1/2$ . Then give the variance of  $\hat{Y}_{\text{new}} = \hat{\beta}_0 + x_{\text{new}} \hat{\beta}_1$  for  $x_{\text{new}} = 3$ .

4. Consider the multiple linear regression model

$$Y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \varepsilon_i$$

for  $i = 1, \dots, 31$ , where  $Y_1, \dots, Y_n$  are volumes of timber harvested from trees,  $x_{11}, \dots, x_{n1}$  are the girths of the trees, and  $x_{12}, \dots, x_{n2}$  are the heights of the trees, and  $\varepsilon_1, \dots, \varepsilon_n$  are assumed to be independent random variables with the Normal( $0, \sigma^2$ ) distribution. Let  $\mathbf{X}$  be the corresponding design matrix and suppose that

$$(\mathbf{X}^T \mathbf{X})^{-1} = \begin{bmatrix} 4.9519 & 0.0287 & -0.0697 \\ 0.0287 & 0.0046 & -0.0012 \\ -0.0697 & -0.0012 & 0.0011 \end{bmatrix}, \quad \hat{\boldsymbol{\beta}} = \begin{bmatrix} -57.9877 \\ 4.7082 \\ 0.3393 \end{bmatrix}, \quad \|\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|_2^2 = 421.9214.$$

In addition  $\sum_{i=1}^n (Y_i - \bar{Y}_n)^2 = 8106.084$ . For the following, you do not need to evaluate your answers.

- (a) Give the value of  $\hat{\sigma}^2$ .
- (b) Give a 95% confidence interval for  $\beta_1$ .
- (c) Suppose you wish to test  $H_0: \beta_2 = 0$  versus  $H_1: \beta_2 \neq 0$  using the  $t$ -test.
  - i. Give the test statistic.
  - ii. Give the rejection rule (give the critical value) under which the test will have size  $\alpha = 0.01$ .
- (d)
  - i. State the null and alternate hypotheses which would be tested by the overall  $F$ -test of significance for this model.
  - ii. Give the value of the test statistic for the overall  $F$ -test of significance,

$$F = \frac{\text{MSM}}{\text{MSE}} = \frac{\text{SSM}/p}{\text{SSE}/(n-p-1)}.$$

- iii. Explain how you would obtain a  $p$ -value for the overall  $F$ -test using the statistic in the previous part. You must give the numerator and denominator degrees of freedom of the relevant  $F$ -distribution.