

STAT 515 Lec 06 slides

Continuous random variables

Karl Gregory

University of South Carolina

These slides are an instructional aid; their sole purpose is to display, during the lecture, definitions, plots, results, etc. which take too much time to write by hand on the blackboard. They are not intended to explain or expound on any material.

Recall: A random variable is a *continuous rv* if its support is an interval¹.

Examples:


- ① Let X be the waiting time until a bus arrives.
- ② Let Y be the time until you drop your new phone.
- ③ Let Z be the body temperature of a randomly selected student.

How do we assign probabilities to the values a continuous rv can take?



- Cannot list vals $x_1, x_2, x_3, \dots$ and assign to them $p_1, p_2, p_3, \dots$ as for discrete.
- Cannot represent the distribution with a table as we have done...

¹or a union of intervals


$X =$ Wait time for hourly departing bus



$$P\left(\frac{1}{4} \leq X\right) = \frac{3}{4}$$



$$P(X = \frac{3}{4}) = 0$$



$$P(a \leq x \leq b) = \int_a^b f(x) dx$$

get Area under f
between a and b .

Probability density function of a continuous rv

The *probability density function (pdf)* of a cont. rv X is the function f satisfying

$$P(a \leq X \leq b) = \int_a^b f(x) dx \quad \text{for all } a \leq b.$$

Note: $\int_a^b f(x) dx = \text{Area between } f \text{ and the horizontal axis on the interval } [a, b].$

Every pdf f has these two properties:

- 1 It is always non-negative, i.e. $f(x) \geq 0$ for all x .
- 2 The total area between f and the horizontal axis is 1, i.e. $\int_{-\infty}^{\infty} f(x) dx = 1$.


If X is an rv with pdf f , then we write $X \sim f$.




ⓐ

We cannot let a pdf take negative values!

Or we'll get negative probs.



$$P(a \leq x \leq b) = \int_a^b f(x) dx$$




Area is > 1

Cannot have this.

$$P(-\infty < x < \infty) = \int_{-\infty}^{\infty} f(x) dx \stackrel{??}{> 1}$$


Discuss: Let X = wait time for hourly arriving bus.

Discuss: Let Y be the time until you drop your new phone. $[0, \infty)$



$$P(98.4 \leq Z \leq 98.8) \stackrel{?}{=} 0.70$$

Discuss: Let Z be the body temperature of a randomly selected student.



let $X \sim f$ (continuous)

$$EX = \int_{-\infty}^{\infty} x f(x) dx = \mu$$

$$Var X = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$$

Expected value and variance of a continuous rv


If $X \sim f(x)$ is a continuous rv, then

- $\mathbb{E}X = \int_{-\infty}^{\infty} x \cdot f(x) dx$
- $\text{Var } X = \int_{-\infty}^{\infty} (x - \mu)^2 \cdot f(x) dx$, where $\mu = \mathbb{E}X$.

Need to use calculus to get these.

$\mathbb{E}X$ is the “balancing point” of the pdf (illustrate).

$\text{Var } X$ describes spread.



$$P(X = x) = 0 .$$

Point probabilities are equal to zero for continuous rvs

For a continuous rv X , for any value c we have

$$P(X = c) = 0,$$

and, as a consequence, for any values a and b such that $a < b$, we have

$$P(a < X < b) = P(a \leq X \leq b) = P(a < X \leq b) = P(a \leq X < b).$$

Exercise: Let X = mpg on next tank of gas.

- ① What is $P(X = 24)$?
- ② What is $P(X = 24.0000000000)$?
- ③ What is $P(24 < X < 25)$ versus $P(24 \leq X < 25)$?
- ④ Say you got $X = 24.2349022301$ last time. Did you observe a 0-prob event??
- ⑤ But how often will that value recur if you fill your tank again and again?



Cumulative distribution function

The *cumulative distribution function (cdf)* F of a rv X is the function F given by

$$F(x) = P(X \leq x) \quad \text{for all } x.$$

If X is an rv with cdf F , then we write $X \sim F$.

We have the following expressions:

$$F(x) = \begin{cases} \int_{-\infty}^x f(t)dt & \text{if } X \text{ is continuous with pdf } f \\ \sum_{t \in \mathcal{X}: t \leq x} p(t) & \text{if } X \text{ is discrete with support } \mathcal{X} \text{ and pmf } p. \end{cases}$$

Exercise: If X = time until you drop your phone, we might have

$$f(x) = \begin{cases} e^{-x}, & x > 0 \\ 0, & x \leq 0 \end{cases} \quad \text{and} \quad F(x) = \begin{cases} 1 - e^{-x}, & x > 0 \\ 0, & x \leq 0. \end{cases}$$

Draw the pdf and cdf.

Exercise: If $X = \#$ jellyfish washed up on the beach, we might have

x	0	1	2	3	4	5	...
$P(X = x)$	0.050	0.149	0.224	0.224	0.168	0.101	...
$F(x) = P(X \leq x)$	0.050	0.199	0.423	0.647	0.815	0.916	...

Draw the pmf and cdf.