

STAT 516 Lec 02

Review of simple linear regression

Karl Gregory

2026-01-21

Hemoglobin versus RBC count example

These data are a subset of the dataset Marcinkevičs et al. (2023)

Some outliers and missing values were removed.

```
link <- url("https://gregorkb.github.io/data/hrbc.csv")
data <- read.csv(link)
head(data)
```

	hem	rbc	sex	age		diag
1	14.8	5.27	female	12.68		appendicitis
2	15.7	5.26	male	14.10	no	appendicitis
3	11.4	3.98	female	14.14	no	appendicitis
4	13.6	4.64	female	16.37	no	appendicitis
5	12.6	4.44	female	11.08		appendicitis
6	12.5	4.96	male	11.05	no	appendicitis

x_i

$$\bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i$$

Hemoglobin level vs red blood cell count for $n = 762$ children.

```
plot(data$hem ~ data$rbc,  
     ylab = "Hemoglobin",  
     xlab = "RBC count")
```


Simple

linear

regression

straight line

one single "predictor" variable.



Simple linear regression

data

Mathematical expression
explaining how the
data arise.

For $(x_1, Y_1), \dots, (x_n, Y_n)$, the simple linear regression model is

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i, \quad i = 1, \dots, n$$


intercept slope error term

where

β = "beta"
 ε = "epsilon"

- ▶ $x_1, \dots, x_n$ are the covariate or predictor values.
- ▶ $Y_1, \dots, Y_n$ are the response values.
- ▶ β_0 and β_1 are the intercept and slope parameters, respectively.
- ▶ $\varepsilon_1, \dots, \varepsilon_n$ are independent $\text{Normal}(0, \sigma^2)$ error terms.
- ▶ σ^2 is the error term variance.

σ^2 "noise" level




3 unknown parameters

β_0 = slope

β_1 = intercept

σ^2 = error term variance





Goals in simple linear regression

$\beta_0, \beta_1, \sigma^2$
— β_1 most interesting

We will learn how to:

1. Estimate the intercept and slope parameters β_0 and β_1 .
2. Estimate the error term variance σ^2 .
3. Perform inference on β_1 .
4. Build a confidence interval for $\beta_0 + \beta_1 x_{\text{new}}$ at any x_{new} .
5. Build a prediction interval for Y at any x_{new} .
6. Decompose the variation in Y into sums of squares.
7. Check whether the model assumptions are satisfied.
8. Identify outliers and understand their effects.

Best line?



Least-squares estimation of slope and intercept

The squared error criterion given by the sum

$$Q(b_0, b_1) = \sum_{i=1}^n (Y_i - (b_0 + b_1 x_i))^2$$

of squared vertical distances of Y_i from the line $y = b_0 + b_1 x$.

We find $Q(b_0, b_1)$ is minimized at $(b_0, b_1) = (\hat{\beta}_0, \hat{\beta}_1)$, where

$$\begin{aligned} \hat{\beta}_0 &= \bar{Y}_n - \hat{\beta}_1 \bar{x}_n \\ \hat{\beta}_1 &= \frac{\sum_{i=1}^n (x_i - \bar{x}_n)(Y_i - \bar{Y}_n)}{\sum_{i=1}^n (x_i - \bar{x}_n)^2} \end{aligned}$$

provided $\sum_{i=1}^n (x_i - \bar{x}_n)^2 > 0$.

The least-squares line or fitted line is the line $y = \hat{\beta}_0 + \hat{\beta}_1 x$.

Pearson's correlation coefficient

Given $(x_1, Y_n), \dots, (x_n, Y_n)$, the quantity

$$\rightarrow r_{xY} = \frac{\sum_{i=1}^n (x_i - \bar{x}_n)(Y_i - \bar{Y}_n)}{\sqrt{\sum_{i=1}^n (x_i - \bar{x}_n)^2 \sum_{i=1}^n (Y_i - \bar{Y}_n)^2}} = \text{cor}(x, Y)$$

is called Pearson's correlation coefficient.

- ▶ Describes strength and direction of *linear* relationships.
- ▶ Must satisfy $r_{xY} \in [-1, 1]$.
- ▶ Values close to zero indicate a weak linear relationship.
- ▶ Is related to $\hat{\beta}_1$ by

$$\rightarrow \hat{\beta}_1 = r_{xY} \frac{S_Y}{S_x} = \text{cor}(x, Y) \frac{\text{sd}(Y)}{\text{sd}(x)}$$

where S_Y and S_x are the sample std devs of the Y and x values.

$$S_x^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}_n)^2$$

Hemoglobin versus RBC count example (cont)

Find the least-squares line on the hemoglobin data.

```
Y <- data$hem
x <- data$rbc
n <- length(Y)
xbar <- mean(x)
Ybar <- mean(Y)


rxY <- cor(x,Y) # Pearson's correlation coefficient
b1hat <- rxY * sd(Y)/sd(x)
b0hat <- Ybar - b1hat * xbar
```

$$\hat{\beta}_0 = 2.99$$

$$\hat{\beta}_1 = 2.16$$

$$y = 2.99 + 2.16x$$

```
plot(data$hem ~ data$rbc, ylab = "Hemoglobin", xlab = "RBC count")
abline(b0hat, b1hat)
```



"Fitted values"
"predicted values"

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

(heights of

RBC count

L-S line at observed covariate values)

Estimating the error term variance

After obtaining $\hat{\beta}_0$ and $\hat{\beta}_1$, define the

- ▶ fitted values as $\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$
- ▶ residuals as $\hat{\varepsilon}_i = Y_i - \hat{Y}_i$

$$\frac{(x_i, y_i)}{n-2}$$

for $i = 1, \dots, n$.

Then an unbiased estimator of σ^2 is given by

$$\hat{\sigma}^2 = \frac{1}{n-2} \sum_{i=1}^n \hat{\varepsilon}_i^2.$$

$$\hat{\sigma} = \sqrt{\frac{1}{n-2} \sum_{i=1}^n \hat{\varepsilon}_i^2}$$

$$= \frac{1}{n-2} \sum_{i=1}^n (\hat{\varepsilon}_i - \bar{\hat{\varepsilon}})^2$$

$= 0.791$ on RBC data.

Confidence interval for the slope parameter

Assume $\varepsilon_1, \dots, \varepsilon_n \stackrel{\text{ind}}{\sim} \text{Normal}(0, \sigma^2)$ and set $S_{xx} = \sum_{i=1}^n (x_i - \bar{x}_n)^2$.

- ▶ The slope estimator $\hat{\beta}_1$ is distributed as

$$\hat{\beta}_1 \sim \text{Normal}(\beta_1, \sigma^2 / S_{xx}).$$

- ▶ “Studentizing” the above gives


$$\frac{\hat{\beta}_1 - \beta_1}{\hat{\sigma} / \sqrt{S_{xx}}} \sim t_{n-2}.$$

- ▶ So a $(1 - \alpha)100\%$ confidence interval for β_1 is

$$\hat{\beta}_1 \pm t_{n-2, \alpha/2} \hat{\sigma} / \sqrt{S_{xx}}.$$

- ▶ We often write $\widehat{\text{s.e.}}(\hat{\beta}_1) = \hat{\sigma} / \sqrt{S_{xx}}$, s.e. for standard error.


Exercise: Justify the CI using the sampling distribution result.



$$\hat{\beta}_1 \sim N\left(\beta_1, \frac{\sigma^2}{S_{xx}}\right)$$

$$\Rightarrow Z = \frac{\hat{\beta}_1 - \beta_1}{\sqrt{\sigma^2 / S_{xx}}} \sim N(0, 1)$$

$$T = \frac{\hat{\beta}_1 - \beta_1}{\sqrt{\hat{\sigma}^2 / S_{xx}}} \sim t_{n-2}$$



$$P\left(-t_{n-2, \alpha/2} < \frac{\hat{\beta}_1 - \beta_1}{\sqrt{\hat{\sigma}^2 / S_{xx}}} < t_{n-2, \alpha/2}\right) = 1 - \alpha$$

$\hat{\sigma} / \sqrt{S_{xx}}$

$$P\left(\hat{\beta}_1 - t_{n-2, \alpha/2} \sqrt{\hat{\sigma}^2 / S_{xx}} < \beta_1 < \hat{\beta}_1 + t_{n-2, \alpha/2} \sqrt{\hat{\sigma}^2 / S_{xx}}\right) = 1 - \alpha$$

PBC data.

$$n = 762$$

$$\alpha = 0.05$$

$$\begin{aligned} t_{762-2, \underbrace{0.05/2}_{0.025}} &= t_{(1-0.025, 762-2)} \\ &= t_{(0.975, 760)} \\ &= 1.963 \end{aligned}$$

$$\hat{\sigma} = 0.491$$

$$\hat{\beta}_1 = 2.163$$

$$S_{xx} = 105.06$$

Hemoglobin versus RBC count example (cont)

Obtain an estimate of the error term variance.


```
Yhat <- b0hat + b1hat * x
ehat <- Y - Yhat
sgsqhat <- sum(ehat^2)/(n-2)
```

We obtain $\hat{\sigma}^2 = 0.626$.

Now construct a 95% confidence interval for β_1 .

```
alpha <- 0.05
Sxx <- sum((x - xbar)^2)
ta2 <- qt(1 - alpha/2, df = n - 2)
lo <- b1hat - ta2 * sqrt(sgsqhat / Sxx)
up <- b1hat + ta2 * sqrt(sgsqhat / Sxx)
```

The 95% CI is (2.011, 2.314).




Tests of hypotheses about the slope

We most often test hypotheses about β_1 of the form

$$\begin{array}{l} H_0: \beta_1 \geq 0 \\ H_1: \beta_1 < 0 \end{array} \quad \text{or} \quad \begin{array}{l} H_0: \beta_1 = 0 \\ H_1: \beta_1 \neq 0 \end{array} \quad \text{or} \quad \begin{array}{l} H_0: \beta_1 \leq 0 \\ H_1: \beta_1 > 0. \end{array}$$

Reject or fail to reject H_0 based on the value of the test statistic

$$T_{\text{stat}} = \frac{(\hat{\beta}_1 - 0)}{\hat{\sigma} / \sqrt{S_{xx}}}$$




Rejection rules for the above at significance level α are

$$T_{\text{stat}} < -t_{n-2,\alpha} \quad \text{or} \quad |T_{\text{stat}}| > t_{n-2,\alpha/2} \quad \text{or} \quad T_{\text{stat}} > t_{n-2,\alpha}.$$

The corresponding p-values are, with $T \sim t_{n-2}$, the probabilities

$$P(T < T_{\text{stat}}) \quad \text{or} \quad 2 \times P(T > |T_{\text{stat}}|) \quad \text{or} \quad P(T > T_{\text{stat}}).$$



2-sided: Reject.

Hemoglobin versus RBC count example (cont)

Test the hypotheses $H_0: \beta_1 = 0$ vs $H_1: \beta_1 \neq 0$ at $\alpha = 0.05$.

```
alpha <- 0.05
Tstat <- b1hat / sqrt(sgsqhat/Sxx)
crit <- qt(1-alpha/2, df = n - 2)
pval <- 2*(1 - pt(abs(Tstat), df = n - 2))
```

We get $T_{\text{stat}} = 28.021$, $t_{n-2, \alpha/2} = 1.963$, and p -value 0; we reject H_0 .

Test the hypotheses $H_0: \beta_1 \leq 2$ vs $H_1: \beta_1 > 2$.

```
Tstat <- (b1hat - 2) / sqrt(sgsqhat/Sxx)
crit <- qt(1-alpha, df = n - 2)
pval <- 1 - pt(Tstat, df = n - 2)
```

We get $T_{\text{stat}} = 2.107$, $t_{n-2, \alpha} = 1.647$, and p -value 0.018; we reject H_0 .

$(x_{\text{new}}, y_{\text{new}})$

↑ will observe later.

A P.I. for new resp value y_{new} .

Want C.I. for height of true line

$$\beta_0 + \beta_1 x_{\text{new}}$$

Expected value / mean response at x_{new} .

$$\hat{y}_{\text{new}} = \hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}}$$

x_{new}

$$\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} \sim N \left(\beta_0 + \beta_1 x_{\text{new}}, \sigma^2 \left[\frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}} \right] \right)$$

Confidence interval for the height of the line

Assume $\varepsilon_1, \dots, \varepsilon_n \stackrel{\text{ind}}{\sim} \text{Normal}(0, \sigma^2)$. Then:

- ▶ The estimator $\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}}$ is distributed as

$$\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} \sim \text{Normal} \left(\beta_0 + \beta_1 x_{\text{new}}, \underbrace{\sigma^2 \left[\frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}} \right]}_{\text{lev}_{\text{new}}} \right).$$

- ▶ “Studentizing” the above gives

$$\frac{\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} - (\beta_0 + \beta_1 x_{\text{new}})}{\hat{\sigma} \sqrt{\frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}}}} \sim \underline{t_{n-2}}.$$

- ▶ So a $(1 - \alpha)100\%$ confidence interval for $\beta_0 + \beta_1 x_{\text{new}}$ is

$$\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} \pm t_{n-2, \alpha/2} \hat{\sigma} \sqrt{\frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}}}.$$

Hemoglobin versus RBC count example (cont)

Give a 95% CI for the mean hemoglobin level of individuals with RBC count 5.5.

```
alpha <- 0.05
xnew <- 5.5
xnew_se <- sqrt(sgsqhat)*sqrt(1/n+(xnew-xbar)^2/Sxx)
ta2 <- qt(1-alpha/2,n-2)
lo <- b0hat + b1hat * xnew - ta2 * xnew_se
up <- b0hat + b1hat * xnew + ta2 * xnew_se
```

We are 95% confident that the mean hemoglobin level of individuals with RBC count 5.5 lies in the interval (14.767, 15.011).

Prediction interval for a new value of the response

Assume $\varepsilon_1, \dots, \varepsilon_n \stackrel{\text{ind}}{\sim} \text{Normal}(0, \sigma^2)$. Then:

- ▶ The new residual $Y_{\text{new}} - \hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}}$ is distributed as

$$Y_{\text{new}} - \hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} \sim \text{Normal} \left(0, \sigma^2 \left[1 + \frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}} \right] \right).$$

- ▶ “Studentizing” the above gives

$$\frac{Y_{\text{new}} - \hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}}}{\hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}}}} \sim t_{n-2}.$$

- ▶ So a $(1 - \alpha)100\%$ prediction interval for Y_{new} is

$$\hat{\beta}_0 + \hat{\beta}_1 x_{\text{new}} \pm t_{n-2, \alpha/2} \hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(x_{\text{new}} - \bar{x}_n)^2}{S_{xx}}}.$$

only difference in the formulas.

Hemoglobin versus RBC count example (cont)

Give a 95% prediction interval for the hemoglobin level when the RBC count is 5.5.

```
alpha <- 0.05
xnew <- 5.5
xnew_pse <- sqrt(sgsqhat)*sqrt(1+1/n+(xnew-xbar)^2/Sxx)
ta2 <- qt(1-alpha/2,n-2)
lo <- b0hat + b1hat * xnew - ta2 * xnew_pse
up <- b0hat + b1hat * xnew + ta2 * xnew_pse
```

We are 95% confident that an *individual* with RBC count 5.5 will have a hemoglobin level in the interval (13.331, 16.447).

Plot confidence and prediction limits over the range of RBC counts.

```
alpha <- 0.05
ta2 <- qt(1-alpha/2,n-2)
xseq <- seq(min(x),max(x),length = 500)


xseq_se <- sqrt(sgsqhat)*sqrt(1/n+(xseq-xbar)^2/Sxx)
loci <- b0hat + b1hat * xseq - ta2 * xseq_se
upci <- b0hat + b1hat * xseq + ta2 * xseq_se

xseq_pse <- sqrt(sgsqhat)*sqrt(1+1/n+(xseq-xbar)^2/Sxx)
lopi <- b0hat + b1hat * xseq - ta2 * xseq_pse
uppi <- b0hat + b1hat * xseq + ta2 * xseq_pse
```

```

plot(Y~x)
abline(b0hat,b1hat)
lines(loci ~ xseq, lty=2); lines(upci ~ xseq, lty=2)
lines(lopi ~ xseq, lty=3); lines(upper ~ xseq, lty=3)

```



The predict() function in R


Can obtain CI for $\beta_0 + \beta_1 x_{\text{new}}$ and PI for Y_{new} with predict() function.

```
lm_out <- lm(Y~x)
xnew <- 5.5
predict(lm_out, newdata = data.frame(x = xnew), int = "conf")
```

	fit	lwr	upr
1	14.88892	14.76725	15.01059

```
predict(lm_out, newdata = data.frame(x = xnew), int = "pred")
```

	fit	lwr	upr
1	14.88892	13.33117	16.44667



$$Q(b_0, b_1) = \sum_{i=1}^n (y_i - (b_0 + b_1 x_i))^2$$

$$Q(\bar{y}_n, 0) = \sum_{i=1}^n (y_i - \bar{y}_n)^2$$

SS_{Tot} = "Total sum of squares"

$$Q(\hat{\beta}_0, \hat{\beta}_1) = \sum_{i=1}^n (y_i - \overbrace{(\hat{\beta}_0 + \hat{\beta}_1 x_i)}^{\text{fitted value}})^2 = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$


$SS_{Error} = \sum_{i=1}^n \hat{\epsilon}_i^2$ = "Error sum of squares"

$$SS_{Reg} = \sum_{i=1}^n (\hat{y}_i - \bar{y}_n)^2 = SS_{Tot} - SS_{Error}$$


Decomposition of variation in $y_1, \dots, y_n$:

$$\underline{SS_{Total}} = \underline{SS_{Reg}} + \underline{SS_{Error}}$$


①



②



③




$$R^2 = \frac{SS_{Reg}}{SS_{Total}}$$

$$0 \leq R^2 \leq 1$$

$$F_{test} = \frac{MS_{Reg}}{MS_{Error}}$$

④



$$\begin{aligned} R^2 &= \frac{SS_{Total} - SS_{Error}}{SS_{Total}} \\ &= 1 - \frac{SS_{Error}}{SS_{Total}} \end{aligned}$$

Sums of squares in simple linear regression

We decompose the variation in $Y_1, \dots, Y_n$ by defining the:

- ▶ Total sum of squares: $SS_{\text{Tot}} = \sum_{i=1}^n (Y_i - \bar{Y}_n)^2$
- ▶ Regression sum of squares: $SS_{\text{Reg}} = \sum_{i=1}^n (\hat{Y}_i - \bar{Y}_n)^2$
- ▶ Error sum of squares: $SS_{\text{Error}} = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$

how different
is the LS
line from
the flat line?

We have $SS_{\text{Tot}} = SS_{\text{Reg}} + SS_{\text{Error}}$.

The coefficient of determination is defined as $R^2 = \frac{SS_{\text{Reg}}}{SS_{\text{Tot}}} = \frac{SS_{\text{Tot}} - SS_{\text{Error}}}{SS_{\text{Tot}}}$


- ▶ $R^2 \in [0, 1]$
- ▶ Proportion of variation in Y "explained" by the covariate x .
- ▶ In simple linear regression we have $R^2 = r_{xY}^2$.

$$= 1 - \frac{SS_{\text{Error}}}{SS_{\text{Tot}}}$$

The mean squares in simple linear regression

The SS, appropriately scaled, follow chi-square distributions:

- ▶ $\frac{SS_{\text{Tot}}}{\sigma^2} \sim \chi_{n-1}^2(\phi_{\text{Total}})$
- ▶ $\frac{SS_{\text{Reg}}}{\sigma^2} \sim \chi_1^2(\phi_{\text{Reg}})$
- ▶ $\frac{SS_{\text{Error}}}{\sigma^2} \sim \chi_{n-2}^2$



The quantities ϕ_{Tot} and ϕ_{Reg} are called noncentrality parameters.

Dividing SS_{Reg} and SS_{Error} by their dfs, we define:

- ▶ Regression mean square: $MS_{\text{Reg}} = \frac{SS_{\text{Reg}}}{1}$

- ▶ Error mean square: $MS_{\text{Error}} = \frac{SS_{\text{Error}}}{n-2}$

$$F_{\text{test}} = \frac{MS_{\text{Reg}}}{MS_{\text{Error}}}$$

The Analysis of Variance (ANOVA) table

$$MS_{Error} = \frac{SS_{Error}}{n-2} = \frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \frac{1}{n-2} \sum_{i=1}^n \hat{\varepsilon}_i^2 = \sigma^2.$$

We often present the SS, df, and MS values in a table like this:

Source	Df	SS	MS	F value	p-value
Regression	1	SS_{Reg}	MS_{Reg}	F_{test}	$P(F > F_{test})$
Error	$n - 2$	SS_{Error}	MS_{Error}		
Total	$n - 1$	SS_{Tot}			

This is an example of an ANOVA table.

Simple linear regression

$$H_0: \beta_1 = 0 \quad \text{vs} \quad H_1: \beta_1 \neq 0.$$

$$T_{test} = \frac{\hat{\beta}_1 - 0}{\hat{\sigma} / \sqrt{S_{xx}}}$$

$$F_{test} = \frac{MS_{Reg}}{MS_{Error}} = \frac{MS_{Reg}}{\hat{\sigma}^2} = T_{test}^2$$

Overall F test

In addition to the SS, df, and MS value, the ANOVA table presents

▶ $F_{\text{stat}} = \frac{\text{MS}_{\text{Reg}}}{\text{MS}_{\text{Error}}}$

▶ $P(F > F_{\text{stat}})$, where this is computed under $F \sim F_{1, n-2}$


These are the test statistic and p-value of the overall F test.

In simple linear regression this p-value is the same as the one for testing $H_0: \beta_1 = 0$ versus $H_1: \beta_1 \neq 0$ with the t test; moreover $F_{\text{stat}} = T_{\text{stat}}^2$.

For what it's worth, one can show $F_{\text{stat}} = \frac{(n-2)r_{xY}^2}{1 - r_{xY}^2}$ in SLR.

We will discuss the overall F test in greater detail later.

$$\frac{\chi^2_{df_1} / df_1}{\chi^2_{df_2} / df_2} \sim F$$



Under $H_0: \beta_1 = 0$ we have


$$F_{test} = \frac{MS_{Reg}}{MS_{Error}} \sim F_{1, n-2}$$

So we reject H_0 if $F_{test} > F_{1, n-2, \alpha}$.
 at significance level α

F., RBC det.

$$n = 762$$

$$\alpha = 0.05$$



$$p\text{-value} = 1 - pt(785.15, 1, 762-2) = 0$$

Building the ANOVA table

```
SST <- sum((Y - Ybar)^2)
SSR <- sum((Yhat - Ybar)^2)
SSE <- sum((Y - Yhat)^2)
MSR <- SSR / 1
MSE <- SSE / (n-2)
Fstat <- MSR / MSE # same as (n-2)*rxY^2/(1 - rxY^2)
pval <- 1 - pf(Fstat,1,n-2)
```

these Dfs should add up to the total.

$$SS_{Error} = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n (y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i))^2$$

	Source	Df	SS	MS	F value	p-value
1	x	1	491.37	491.37	785.15	0
n-2	Error	760	475.63	0.63		$\frac{491.37}{0.63}$
	Total	761	967			

$$n = 762$$

$$n-1$$

$$SS_{Total} = \sum_{i=1}^n (y_i - \bar{y})^2$$

The `lm()`, `summary()`, and `anova()` functions in R

```
lm_out <- lm(Y~x)
lm_out
```

Call:

```
lm(formula = Y ~ x)
```

Coefficients:

(Intercept)	x
2.994	2.163

```
summary(lm_out)
```

Call:

```
lm(formula = Y ~ x)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.9702	-0.4232	0.0074	0.4645	2.3791

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	2.99447	0.37065	8.079	2.56e-15 ***
x	2.16263	0.07718	28.021	< 2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.7911 on 760 degrees of freedom

Multiple R-squared: 0.5081, Adjusted R-squared: 0.5075

F-statistic: 785.1 on 1 and 760 DF, p-value: < 2.2e-16

```
anova(lm_out)
```

Analysis of Variance Table

Response: Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
x	1	491.37	491.37	785.15	< 2.2e-16 ***
Residuals	760	475.63	0.63		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Checking model assumptions


$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$


independent
 $\varepsilon_i \sim N(0, \sigma^2)$
 $i = 1, \dots, n.$

Validity of the foregoing analyses depends on these assumptions:


1. The responses are normally distributed around the regression line (Check QQ plot of residuals). *If n is large this doesn't matter.*
2. The response has the same variance for all values of the covariate (Check residuals vs fitted values plot).
3. The covariate and the response are linearly related (Check residuals vs fitted values plot).
4. The response values are independent of each other (No way to check; must trust experimental design).

When to see random scatter
in resid. vs fitted values plot.






① Get residuals, Normal Q-Q plot.




Generating diagnostic plots from `lm()` with `plot()`

```
plot(lm_out, which = 2)
```



```
plot(lm_out, which = 1)
```



The abalone...



Photo on the left by Sharktopus - Own work, CC BY-SA 3.0,
<https://commons.wikimedia.org/w/index.php?curid=14082271>

Abalone data example


Predict shucked weight of an abalone by its length.

```
csv <- url("https://people.stat.sc.edu/gregorkb/data/abalone.csv")
abalone <- read.csv(csv,col.names = c("Sex",
                                      "Length",
                                      "Diameter",
                                      "Height",
                                      "Whole_Wt",
                                      "Shucked_Wt",
                                      "Viscera_Wt",
                                      "Shell_Wt",
                                      "Rings"))

Y <- abalone$Shucked_Wt
x <- abalone$Length
n <- length(Y)
```


There are $n = 4176$ records. Data come from Nash and Ford (1995).

```
lm1 <- lm(Y~x)
plot(Y~x)
abline(lm1)
```




Try transforming x:

```
x3 <- x**3  
lm2 <- lm(Y ~ x3)  
plot(Y ~ x3); abline(lm2)
```



```
plot(lm2, which = 1)
```



Maybe:

$$\% \text{ change in weight} = \beta_0 + \beta_1 \% \text{ change in length.}$$


Could transform both Y and x:

```
logY <- log(Y); logx <- log(x)
lm3 <- lm(logY ~ logx)
plot(logY~logx); abline(lm3)
```


$$\frac{d}{dy} \log y = \frac{1}{y}$$

$$\frac{d}{dx} \log x = \frac{1}{x}$$


$$\log y = \beta_0 + \beta_1 \log x \Rightarrow \frac{dy}{y} = \beta_1 \frac{dx}{x}$$



```
plot(lm3, which = 2)
```



```
plot(lm3, which = 1)
```



Transforming variables to obtain a linear relationship

Take care how to interpret β_1 after transforming the data.

Example: Log transforming x and Y gives β_1 a %-change interpretation:

$$\log y = \beta_0 + \beta_1 \log x \iff \frac{d \log y}{dx} = \beta_1 \frac{1}{x} \iff \frac{dy}{y} = \beta_1 \frac{dx}{x}$$

Abalone data example (cont)

We must back-transform prediction intervals if we have transformed Y .

```
xnew <- 0.5
newdata <- data.frame( logx = log(xnew))
pi_logY <- predict(lm3,newdata = newdata, int = "pred")
pi_logY
```

	fit	lwr	upr
1	-1.335636	-1.724831	-0.946441


```
pi <- exp(pi_logY)
pi
```

	fit	lwr	upr
1	0.2629909	0.1782032	0.3881199

```

plot(Y~x,col="gray"); logx <- seq(min(logx),max(logx),length=500)
newdata <- data.frame(logx = logx)
logy_hat <- predict(lm3,newdata = newdata,int = "pred")
lines(exp(logy_hat[,1]) ~ exp(logx), col = "red")
lines(exp(logy_hat[,2]) ~ exp(logx), col = "red", lty = 3,lwd=1.5)
lines(exp(logy_hat[,3]) ~ exp(logx), col = "red", lty = 3,lwd=1.5)

```




Outliers in simple linear regression

Outlying data points can have a large influence on the estimated regression function.


Let's generate some data and then add an outlier:

```
n <- 20
b0 <- 1
b1 <- -1/2
sg <- .2
x0 <- runif(n,0,5)
e <- rnorm(n,0,sg)
Y0 <- b0 + b1 * x0 + e
x <- c(x0,.3)
Y <- c(Y0,-1.3)
```

```
plot(Y~x);points(Y[n+1]~x[n+1], col = "red")
abline(lm(Y0~x0))
abline(lm(Y~x), col = "red")
```



The red data point appears to exert an undue influence over the fit.



leverage:
$$\text{lev}_i = \frac{1}{n} + \frac{(x_i - \bar{x}_n)^2}{S_{xx}}$$

Leverage and Cook's distance

The leverage of a point (x_i, Y_i) among $(x_1, Y_1), \dots, (x_n, Y_n)$ is

$$\text{lev}_i = \frac{1}{n} + \frac{(x_i - \bar{x}_n)^2}{S_{xx}}$$

Leverage only shows outlying-ness in the x direction.

Cook's Distance measures how much each data point changes the fit:

$$D_i = \frac{1}{2\hat{\sigma}^2} \sum_{j=1}^n (\hat{Y}_j - \hat{Y}_{j(i)})^2 = \frac{\hat{e}_i^2}{2\hat{\sigma}^2} \frac{\text{lev}_i}{(1 - \text{lev}_i)^2} \quad \text{for } i = 1, \dots, n,$$


after removing data point i.

where $\hat{Y}_{j(i)}$ is the j th fitted value from the model fitted without obs i .

height of L-S line on full data at $x = x_j$

Make a plot of the Cook's distances

```
plot(lm(Y~x),which = 4)
```



Code to compute Cook's distances

```
n <- length(Y)
xbar <- mean(x)
Ybar <- mean(Y)
rxY <- cor(x,Y) # Pearson's correlation coefficient
b1hat <- rxY * sd(Y)/sd(x)
b0hat <- Ybar - b1hat * xbar
Sxx <- sum((x - xbar)^2)
lev <- 1/n + (x - xbar)^2/Sxx
Yhat <- b0hat + b1hat * x
ehat <- Y - Yhat
sgsqhat <- sum(ehat^2)/(n-2)

cooksD <- ehat^2 / (2*sgsqhat) * lev / (1 - lev)^2
```

Two-sample t-test by simple linear regression

Let $Y_{ij} \stackrel{\text{ind}}{\sim} \text{Normal}(\mu_i, \sigma^2)$, $j = 1, \dots, n_i$, $i = 1, 2$ and consider

$$H_0: \mu_2 - \mu_1 = 0 \text{ versus } H_1: \mu_2 - \mu_1 \neq 0.$$

The (equal-variances) two-sample t-test uses the test statistic

$$T_{\text{stat}} = \frac{\bar{Y}_2 - \bar{Y}_1}{S_{\text{pooled}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}},$$

where $\bar{y}_i = n_i^{-1} \sum_{j=1}^{n_i} Y_{ij}$, $i = 1, 2$ and

$$S_{\text{pooled}}^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}, \quad S_i = \frac{1}{n_i - 1} \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2.$$

We reject H_0 at significance level α if $|T_{\text{stat}}| > t_{n-2, \alpha/2}$.

Appendicitis example

Look again at the data from Marcinkevičs et al. (2023).


```
link <- url("https://people.stat.sc.edu/gregorkb/data/hrbc.csv")
data <- read.csv(link)
head(data)
```

	hem	rbc	sex	age	diag
1	14.8	5.27	female	12.68	appendicitis
2	15.7	5.26	male	14.10	no appendicitis
3	11.4	3.98	female	14.14	no appendicitis
4	13.6	4.64	female	16.37	no appendicitis
5	12.6	4.44	female	11.08	appendicitis
6	12.5	4.96	male	11.05	no appendicitis

Is the mean hemaglobin level the same in children with and without appendicitis (ignoring rbc, age, and sex)?

Appendicitis example (cont)

```
boxplot(data$hem ~ data$diag)
```



Appendicitis example (cont)

```
t.test(data$hem ~ data$diag, var.equal = TRUE)
```

Two Sample t-test

data: data\$hem by data\$diag

t = -0.49212, df = 760, p-value = 0.6228

alternative hypothesis: true difference in means between group appendicitis and

95 percent confidence interval:

-0.2038964 0.1221585

sample estimates:

mean in group appendicitis	mean in group no appendicitis
----------------------------	-------------------------------

13.33229	13.37316
----------	----------

Appendicitis example (cont)

Let the Y_i be the hemaglobin values and define an indicator variable as

$$x_i = \begin{cases} 0 & \text{if no appendicitis} \\ 1 & \text{if appendicitis} \end{cases} \quad \text{for } i = 1, \dots, n.$$

Then in the SLR model $Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$ we have

- ▶ $\beta_0 = \mu_{\text{no app}}$
- ▶ $\beta_0 + \beta_1 = \mu_{\text{app}}$
- ▶ $\beta_1 = \mu_{\text{app}} - \mu_{\text{no app}}$

The t test in the simple linear regression setup of

$$H_0: \beta_1 = 0 \text{ versus } H_1: \beta_1 \neq 0$$

will give the same p value as the equal-variances two-sample t test of

$$H_0: \mu_{\text{app}} - \mu_{\text{no app}} = 0 \text{ versus } H_1: \mu_{\text{app}} - \mu_{\text{no app}} \neq 0. \quad \text{Cool!}$$

Exercise: Show that in the above setup we have

$$\frac{\bar{Y}_2 - \bar{Y}_1}{S_{\text{pooled}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} = \frac{\hat{\beta}_1}{\hat{\sigma} / \sqrt{S_{xx}}}.$$

Do it in steps, showing:

1. $\hat{\beta}_0 = \bar{Y}_1$
2. $\hat{\beta}_1 = \bar{Y}_2 - \bar{Y}_1$
3. $\hat{\sigma} = S_{\text{pooled}}$
4. $1 / \sqrt{S_{xx}} = \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$

Appendicitis example (cont)

Prepare the data:

```
Y <- data$hem  
x <- as.numeric(data$diag == "appendicitis")  
head(cbind(Y,x))
```

	Y	x
[1,]	14.8	1
[2,]	15.7	0
[3,]	11.4	0
[4,]	13.6	0
[5,]	12.6	1
[6,]	12.5	0

```
summary(lm(Y~x))
```

Call:

```
lm(formula = Y ~ x)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.1323	-0.7323	-0.0323	0.6677	4.1677

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	13.37316	0.06375	209.782	<2e-16 ***
x	-0.04087	0.08305	-0.492	0.623

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.128 on 760 degrees of freedom

Multiple R-squared: 0.0003186, Adjusted R-squared: -0.0009968

F-statistic: 0.2422 on 1 and 760 DF, p-value: 0.6228

Automatic if we designate the predictor as a “factor” (but watch sign!).

```
x <- as.factor(data$diag)
summary(lm(data$hemo ~ x))
```

Call:

```
lm(formula = data$hemo ~ x)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.1323	-0.7323	-0.0323	0.6677	4.1677

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	13.33229	0.05322	250.490	<2e-16 ***
xno appendicitis	0.04087	0.08305	0.492	0.623

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.128 on 760 degrees of freedom


Multiple R-squared: 0.0003186, Adjusted R-squared: -0.0009968

F-statistic: 0.2422 on 1 and 760 DF, p-value: 0.6228

The importance (or unimportance) of Normal error terms


Generate some data with a right-skewed error term distribution:

```
b0 <- 0; b1 <- 1; n <- 500; x <- rnorm(n); e <- rgamma(n, shape = 3/2, scale = 2/3) - 1  
Y <- b0 + b1*x + e  
plot(Y~x); lm_out <- lm(Y~x); abline(lm_out, col = "red")
```



Check the Normal quantile-quantile plot of the residuals.

```
plot(lm_out, which = 2)
```



Generate a large number of such data sets and obtain $\hat{\beta}_1$ for each one.

```
S <- 300
b1hat <- numeric(S)

for(s in 1:S){


  x <- rnorm(n)
  e <- rgamma(n, shape = 3/2, scale = 2/3) - 1
  Y <- b0 + b1*x + e
  lm_out <- lm(Y~x)
  b1hat[s] <- coef(lm_out)[2]

}
```

Check if the $\hat{\beta}_1$ values have a Normal distribution.

```
qqnorm(scale(b1hat))  
abline(0,1)
```

Normal Q–Q Plot



References

- Marcinkevičs, Ričards, Patricia Reis Wolfertstetter, Ugne Klimiene, Ece Ozkan, Kieran Chin-Cheong, Alyssia Paschke, Julia Zerres, et al. 2023. “Regensburg Pediatric Appendicitis Dataset.” Zenodo. <https://doi.org/10.5281/zenodo.7711412>.
- Nash, Sellers, Warwick, and Wes Ford. 1995. “Abalone.” UCI Machine Learning Repository. <https://doi.org/10.24432/C55C7W>.