

STAT 214 HW 02 SOLUTIONS

2

Four treatments will be compared in an experiment in which four subjects are assigned to each treatment according to a Latin Square design.

There are two blocking variables - row and column blocking variables - with four levels each.

The block and treatment arrangement will follow the diagram

	B_1	B_2	B_3	B_4
A_1	1	2	3	4
A_2	2	1	4	3
A_3	3	4	2	1
A_4	4	3	1	2

where A_1, A_2, A_3, A_4 and B_1, B_2, B_3, B_4 are block effects, and the numbers in the cell indicate what treatment is applied at the block combinations.

The resulting data will be analyzed assuming the linear model

$$Y_{ijk} = \mu + A_i + B_j + d_k + \varepsilon_{ijk}, \quad i, j, k = 1, 2, 3, 4,$$

where

- μ is a mean
- d_1, \dots, d_4 are treatment effects
- A_i are indep. $N(0, \sigma_A^2)$
- B_i are indep. $N(0, \sigma_B^2)$
- ε_{ijk} are indep. $N(0, \sigma_\varepsilon^2)$
- A_i, B_i , and ε_{ijk} are independent.

Write the linear model in matrix notation

$$\underline{y} = \underline{X}\underline{b} + \underline{Z}\underline{u} + \underline{\varepsilon},$$

where \underline{b} contains fixed parameters and \underline{u} contains random effects.

Write out the entries of each vector and matrix.

Solution:

[illegible]

[2] For a matrix $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ with A and D invertible, verify that

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} + A^{-1}BE^{-1}CA^{-1} & -A^{-1}BE^{-1} \\ -E^{-1}CA^{-1} & E^{-1} \end{bmatrix},$$

where $E = D - CA^{-1}B$

Solution:

$$\begin{bmatrix} A^{-1} + A^{-1}BE^{-1}CA^{-1} & -A^{-1}BE^{-1} \\ -E^{-1}CA^{-1} & E^{-1} \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}$$

$$= \begin{bmatrix} I + A^{-1}BE^{-1}C - A^{-1}BE^{-1}C & A^{-1}B + A^{-1}BE^{-1}CA^{-1}B - A^{-1}BE^{-1}D \\ -E^{-1}C + E^{-1}C & -E^{-1}CA^{-1}B + E^{-1}D \end{bmatrix}$$

Use $D = E + CA^{-1}B$

$$= \begin{bmatrix} I & A^{-1}B + A^{-1}BE^{-1}CA^{-1}B - A^{-1}BE^{-1}(E + CA^{-1}B) \\ 0 & -E^{-1}CA^{-1}B + E^{-1}(E + CA^{-1}B) \end{bmatrix}$$

$$= \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}.$$

4] Let X be an $n \times p$ matrix.

Describe the change in X when it is premultiplied by $\left(I_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T\right)$.

Solution: We have

$$\begin{aligned} \left(I_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T\right) X &= \left(I_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T\right) [x_1 \cdots x_p] \\ &= \left[x_1 - \left(\frac{1}{n} \sum_{i=1}^n x_{1i}\right) \mathbf{1}_n \cdots x_p - \left(\frac{1}{n} \sum_{i=1}^n x_{pi}\right) \mathbf{1}_n \right], \end{aligned}$$

So premultiplication of X by $\left(I_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T\right)$ centers the columns of X so that they have mean zero.

5 Characterize the solution set of $A\vec{x} = \vec{b}$ (provided the system of equations is consistent), where

$$A = \begin{bmatrix} 3 & 5 & -2 \\ -3 & -2 & -1 \\ 6 & 1 & 5 \end{bmatrix} \quad \text{and} \quad \vec{b} = \begin{bmatrix} -7 \\ 1 \\ 4 \end{bmatrix}.$$

Solution: Row-reduce the augmented matrix:

$$[A \quad \vec{b}] = \begin{bmatrix} 3 & 5 & -2 & -7 \\ -3 & -2 & -1 & 1 \\ 6 & 1 & 5 & 4 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & 5 & -2 & -7 \\ 0 & 3 & -3 & -6 \\ 0 & -9 & 9 & 18 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & 5 & -2 & -7 \\ 0 & 3 & -3 & -6 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & 5 & -2 & -7 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & 0 & 3 & 3 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} x_1 + x_3 &= 1 & x_1 &= 1 - x_3 \\ x_2 - x_3 &= -2 & \Leftrightarrow x_2 &= -2 + x_3 \\ x_3 &\text{ free} & x_3 &= x_3 \end{aligned}$$

So the solution set is given by

$$\left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}, \quad x_3 \in \mathbb{R} \right\}$$

6 Show that if two nonzero vectors \underline{v}_1 and \underline{v}_2 are orthogonal, then $\{\underline{v}_1, \underline{v}_2\}$ is linearly independent.

Solution: Orthogonality of \underline{v}_1 and \underline{v}_2 means $\underline{v}_1 \cdot \underline{v}_2 = 0$.

Suppose $\underline{v}_1 c_1 + \underline{v}_2 c_2 = 0$.

Pre-multiplying by \underline{v}_j^T gives $\underline{v}_j^T \underline{v}_j c_j = 0$ for $j=1,2$.

Since \underline{v}_1 and \underline{v}_2 are nonzero, we must have $c_1 = c_2 = 0$.

Therefore \underline{v}_1 and \underline{v}_2 are linearly independent.

7 Let $\{\underline{v}_1, \underline{v}_2\}$ be a set of linearly independent vectors in \mathbb{R}^n and let

$$\underline{u}_1 = \underline{v}_1 \quad \text{and} \quad \underline{u}_2 = \underline{v}_2 - \left(\frac{\underline{v}_2 \cdot \underline{v}_1}{\underline{v}_1 \cdot \underline{v}_1} \right) \underline{v}_1.$$

Show that $\{\underline{u}_1, \underline{u}_2\}$ is linearly independent. Hint: Show that \underline{u}_1 and \underline{u}_2 are orthogonal.

Solution: We have

$$\begin{aligned} \underline{u}_1 \cdot \underline{u}_2 &= \underline{v}_1 \cdot \left[\underline{v}_2 - \left(\frac{\underline{v}_2 \cdot \underline{v}_1}{\underline{v}_1 \cdot \underline{v}_1} \right) \underline{v}_1 \right] \\ &= \underline{v}_1 \cdot \underline{v}_2 - \underline{v}_1 \cdot \underline{v}_1 \left(\frac{\underline{v}_2 \cdot \underline{v}_1}{\underline{v}_1 \cdot \underline{v}_1} \right) \\ &= 0. \end{aligned}$$

Since \underline{u}_1 and \underline{u}_2 are orthogonal, $\{\underline{u}_1, \underline{u}_2\}$ is linearly independent.