

## STAT 720 sp 2019 midterm

assigned: Wednesday, March 20th, 2019

*due on Friday, March 22nd, 2019 at 5:00 pm in my mailbox or to me in my office.*

This is a take-home exam. You may use all the course notes and any books. Do not work together with any classmates or with any other person.

1. Consider the MA(2) time series  $\{X_t, t \in \mathbb{Z}\}$  defined by

$$X_t = \mu + Z_t + \theta_1 Z_{t-1} + \theta_2 Z_{t-2}, \quad \text{for all } t \in \mathbb{Z}, \quad (1)$$

where  $\{Z_t, t \in \mathbb{Z}\} \sim \text{IID}(0, \sigma^2)$ .

- (a) Find the autocovariance function  $\gamma(\cdot)$  of the time series  $\{X_t, t \in \mathbb{Z}\}$ .
- (b) Find  $2\pi f(0)$ , where  $f(\cdot)$  is the spectral density of the time series  $\{X_t, t \in \mathbb{Z}\}$ .
- (c) Verify that  $2\pi f(0) = \sum_{h=-\infty}^{\infty} \gamma(h)$ .
- (d) Give the form of an asymptotic 95% confidence interval for  $\mu$  based on  $X_1, \dots, X_n$ , assuming  $\theta_1$ ,  $\theta_2$ , and  $\sigma^2$  are known.
- (e) Give finite values of the parameters  $\theta_1$  and  $\theta_2$ , if they exist, for which the time series  $\{X_t, t \in \mathbb{Z}\}$  is
  - i. invertible.
  - ii. non-invertible.
  - iii. non-stationary.
  - iv.  $m$ -dependent with  $m = 1$ .

For each case explain your answer.

- (f) Let  $f(\cdot)$  be the spectral density of a stationary time series with autocovariance function  $\gamma(\cdot)$ . Show that  $\int_{-\pi}^{\pi} f(\lambda) d\lambda = \gamma(0)$ .
  - (g) For the time series  $\{X_t, t \in \mathbb{Z}\}$  in (1), give  $\int_{-\pi}^{\pi} f(\lambda) d\lambda$  in terms of  $\theta_1$ ,  $\theta_2$ , and  $\sigma^2$ .
  - (h) Assuming that  $\mu$ ,  $\theta_1$ ,  $\theta_2$ , and  $\sigma^2$  are known, give the one-step-ahead predictor  $\hat{X}_1$  of  $X_1$  based on  $X_1, \dots, X_n$  and give the mean squared error of prediction (MSEP) associated with the predictor.
  - (i) Assuming that  $\mu$ ,  $\theta_1$ ,  $\theta_2$ , and  $\sigma^2$  are known, give the  $h$ -step-ahead predictors  $\hat{X}_{n+h}$  of  $X_{n+h}$  for  $h > 2$  based on  $X_1, \dots, X_n$  and give the MSEP associated with the predictors.
  - (j) Describe in words how you would generate a realization of length 100 of the time series  $\{X_t, t \in \mathbb{Z}\}$ , where  $\{Z_t, t \in \mathbb{Z}\}$  are independent  $\text{Normal}(0, \sigma^2)$  random variables.
2. For each of the following time series, indicate which statements among the following four statements, if any, must be true.

- |                               |                                     |
|-------------------------------|-------------------------------------|
| (I) it is strictly stationary | (III) it is not strictly stationary |
| (II) it is stationary         | (IV) it is not stationary           |

- (a) Let  $\{Y_t, t \in \mathbb{Z}\}$  be a time series such that the covariance matrix containing the covariances among  $Y_1, \dots, Y_5$  is given by

$$(\text{Cov}(Y_i, Y_j))_{1 \leq i, j \leq 5} = \begin{bmatrix} 1 & 0.9 & 0 & 0 & 0 \\ 0.9 & 1 & 0.9 & 0 & 0 \\ 0 & 0.9 & 1 & 0.8 & 0 \\ 0 & 0 & 0.8 & 1 & 0.8 \\ 0 & 0 & 0 & 0.8 & 1 \end{bmatrix}$$


- (b) Let  $\{Y_t, t \in \mathbb{Z}\}$  be a time series such that  $\mathbb{E}Y_t = \mu$  for all  $t \in \mathbb{Z}$  and  $\text{Cov}(Y_i, Y_j) = (0.9)^{|i-j|}$  for  $i, j \in \mathbb{Z}$ .
- (c) Let  $\{Y_t, t \in \mathbb{Z}\}$  be a time series such that

$$Y_t = (3/4)Y_{t-1} + (1/4)Y_{t-2} + Z_t, \quad t \in \mathbb{Z},$$

where  $\{Z_t, t \in \mathbb{Z}\}$  are independent  $\text{Normal}(0, \sigma^2)$  random variables.

- (d) Let  $\{Y_t, t \in \mathbb{Z}\}$  be a time series such that  $Y_t \sim \text{Poisson}(\lambda)$  for all  $t \in \mathbb{Z}$ .
- (e) Let  $\{Y_t, t \in \mathbb{Z}\}$  be a time series such that  $Y_t \sim \text{Normal}(0, \sigma^2)$  for all  $t \in \mathbb{Z}$  and  $\text{Cov}(Y_i, Y_j) = \mathbf{1}(|i - j| > 0)$  for  $i, j \in \mathbb{Z}$ , where  $\mathbf{1}(\cdot)$  is the indicator function.

3. Suppose you play a game of Monopoly and you record after each turn the number of spaces you must move in order to reach the GO tile, resulting in a time series  $\{X_t, t = 1, 2, \dots\}$ . There are 40 tiles on the Monopoly board; you begin on the GO tile and advance by moving a number of tiles equal to the sum of two dice rolls (the dice are six-sided). Your observed time series might look like this after 100 turns:



- (a) Find  $\mathbb{E}X_1$ , where  $X_1$  is the number of spaces you must move to reach the GO tile after your first turn.
- (b) Find  $\mathbb{E}X_2$ , where  $X_2$  is the number of spaces you must move to reach the GO tile after your second turn.

- (c) Suggest a value for  $\lim_{n \rightarrow \infty} \mathbb{E}X_t$  and argue why it is correct.
- (d) Describe a transformation of the time series  $\{X_t, t = 1, 2, \dots\}$  which would result in a strictly stationary time series.
4. Let  $U$  and  $V$  be random variables such that  $\text{Var } U = \sigma_U^2$ ,  $\text{Var } V = \sigma_V^2$ , and  $\text{Cov}(U, V) = \sigma_{UV}$ . You wish to predict the value of  $U$  using the value of  $V$  with a predictor of the form

$$\hat{U} = a_0 + a_1 V$$

for some  $a_0, a_1 \in \mathbb{R}$ .

- (a) Find the values  $a_0$  and  $a_1$  which minimize the MSEP of the predictor  $\hat{U}$ ; that is, find  $a_0$  and  $a_1$  which minimize  $\mathbb{E}(U - \hat{U})^2$ .
- (b) Give the MSEP of  $\hat{U}$  under the choices of  $a_0$  and  $a_1$  from part (a).
5. Let  $V_1, \dots, V_n, V_{n+1}$  be random variables with zero mean such that

$$\text{Cov}(V_i, V_j) = \begin{cases} 1, & i = j \\ 1/4, & |i - j| \leq 3 \\ 0, & |i - j| > 3 \end{cases}$$

Consider a predictor  $\hat{V}_{n+1}$  of  $V_{n+1}$  based on  $V_1, \dots, V_n$  which is of the form

$$\hat{V}_{n+1} = \sum_{i=1}^n a_i V_{n+1-i}.$$

For  $n = 2$ , find the values of  $a_1, \dots, a_n$  which minimize the MSEP of the predictor  $\hat{V}_{n+1}$ .

6. Consider the causal ARMA(2, 1) time series given by

$$X_t - \phi_1 X_{t-1} - \phi_2 X_{t-2} = Z_t + \theta_1 Z_{t-1}, \quad t \in \mathbb{Z},$$

where  $\{Z_t, t \in \mathbb{Z}\}$  is  $\text{WN}(0, \sigma^2)$ . Let  $\{\psi_j, j = 0, 1, \dots\}$  be the coefficients such that we may write

$$X_t = \sum_{j=0}^{\infty} \psi_j Z_{t-j}, \quad t \in \mathbb{Z}.$$

Give  $\psi_0, \psi_1, \psi_2$ , and  $\psi_3$ , in terms of  $\phi_1, \phi_2$ , and  $\theta_1$ .