

Formula Sheet – Final Exam – SCCC 312A
Classical (Wald) $1 - \alpha$ CI for p :

$$\left(\hat{p} - z(\alpha/2) \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}, \hat{p} + z(\alpha/2) \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right)$$

$1 - \alpha$ CI for μ :

$$\left(\bar{x} - t(n - 1, \alpha/2) \frac{s}{\sqrt{n}}, \bar{x} + t(n - 1, \alpha/2) \frac{s}{\sqrt{n}} \right)$$

Test statistic (hypothesis test about μ):

$$t^* = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

CI and test statistics for μ_d in paired-sample t -test: Same as above but with \bar{x}_d and s_d .

Test statistic (hypothesis test about p):

$$z^* = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

$1 - \alpha$ CI for $\mu_1 - \mu_2$:

$$\left((\bar{x}_1 - \bar{x}_2) - t(df, \alpha/2) \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}, (\bar{x}_1 - \bar{x}_2) + t(df, \alpha/2) \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} \right)$$

Test statistic (comparing two means, independent samples):

$$t^* = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

$1 - \alpha$ CI for $p_1 - p_2$:

$$\left((\hat{p}_1 - \hat{p}_2) - z(\alpha/2) \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}, (\hat{p}_1 - \hat{p}_2) + z(\alpha/2) \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} \right)$$

Test statistic (comparing two proportions):

$$z^* = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

where \hat{p} is the pooled sample proportion.