

Stat 512 — Take home exam IV (due on July 29th)

1. In a study of the relationship between birth order and college success, an investigator found that 126 in a sample of 180 college graduates were firstborn or only children; in a sample of 100 nongraduates of comparable age and socioeconomic background, the number of firstborn or only children was 54. Estimate the difference in proportions of firstborn or only children for the two populations from which these samples were drawn using 90% confidence coefficient. Is there sufficient evidence to conclude the proportion of firstborn or children for college graduates is higher than that of nongraduates? (10 pts)
2. A small amount of the trace element selenium, from 50 to 200 micrograms per day, is considered essential to good health. Suppose that independent random samples of $n_1 = n_2 = 30$ adults were selected from two regions of the United States, and a days intake of selenium, from both liquids and solids, was recorded for each person. The mean and standard deviation of the selenium daily intakes for the 30 adults from region 1 were $\bar{Y}_1 = 165.1$ and $S_1 = 23$, respectively. The corresponding statistics for the 30 adults from region 2 were $\bar{Y}_2 = 138$ and $S_2 = 22$. Find a 90% confidence interval for the difference in the mean selenium intake for the two groups. (10 pts)
3. Organic chemists often purify organic compounds by a method known as fractional crystallization. An experimenter wanted to prepare and purify 4.85g of aniline. Ten 4.85-gram specimens of aniline were prepared and purified to produce acetanilide. The following dry yields were obtained:

3.85, 3.88, 3.90, 3.62, 3.72, 3.80, 3.85, 3.36, 4.01, 3.82

Construct a 95% CI for the mean number of grams of acetanilide that can be recovered from 4.85 grams of aniline. (10 pts) (Hint: 1. you need to use software to calculate sample mean, variance and t critical value. 2. Be careful about the formula you choose. Is it a small sample or large sample situation?)

4. Let Y has pdf:

$$f_Y(y) = \frac{2(\theta - y)}{\theta^2}, \quad y \in (0, \theta)$$

- a. Find out the CDF of Y . (10 pts)
- b. Show that $\frac{Y}{\theta}$ is a pivotal quantity. (10 pts)
- c. Use the pivotal quantity from part (b) to find a 90% upper confidence limit for θ . (10 pts)

5. The distribution function for a power family distribution is given by:

$$F_Y(y) = \left(\frac{y}{\theta}\right)^\alpha, \quad y \in [0, \theta]$$

where $\alpha, \theta > 0$. Assume that a sample of size n is taken from a population with a power family distribution and that $\alpha = c$ where $c > 0$ is known.

a. Show that the distribution function of $Y_{(n)} = \max\{Y_1, \dots, Y_n\}$ is given by: (10 pts)

$$F_{Y_{(n)}}(y) = \left(\frac{y}{\theta}\right)^{nc}, \quad y \in [0, \theta]$$

b. Show that $\frac{Y_{(n)}}{\theta}$ is a pivotal quantity and that for $0 < k < 1$ (10 pts)

$$P\left(k < \frac{Y_{(n)}}{\theta} \leq 1\right) = 1 - k^{cn}$$

c. Suppose that $n = 5$ and $\alpha = c = 2.4$, then use the result from part (b) to find k such that (10 pts)

$$P\left(k < \frac{Y_{(5)}}{\theta} \leq 1\right) = 0.95$$

and give a 95% CI for θ .

6. Let X_1, \dots, X_{10} are iid with pdf $f(x|\mu) = e^{-(x-\mu)}$, where $x > \mu$ (This is called a shifted exponential distribution). Find a $100(1 - \alpha)\%$ CI for μ using the pivotal method. (Hint: Consider a function of $Y_{(1)} = \min(X_1, \dots, X_{10})$). (10 pts)